

SUPPLEMENTAL LABELING



PYRIFLUQUINAZON GROUP 9B INSECTICIDE

EPA Reg. No. 71711-37

For Use On: Citrus Fruits (Crop Group 10-10)

Use Pattern: Aerial Application

This supplemental labeling expires 07/31/2023 and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application. Read the label affixed to the container for PQZ[®] Insecticide before applying. Use of PQZ Insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for PQZ Insecticide. New use directions appear on this supplemental labeling that may be different from those that appear on the container label.

BUFFER ZONES

Vegetative Filter (Buffer) Strip

All crops: 15-foot vegetative filter (buffer) strip

Construct and maintain the vegetative filter (buffer) strip of grass or other permanent vegetation between field edge and down gradient aquatic habitat (including but not limited to lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds).

Only apply products containing pyrifluquinazon onto fields where a maintained vegetative filter (buffer) strip of at least 15 feet exists between the field edge and down gradient aquatic habitat. For guidance, refer to the following publication for information on constructing and maintaining effective buffers: Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services. USDA, https://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs144p2_030970.pdf

Buffer Zone for Aerial Application

Do not apply within 150 feet of aquatic habitats (including but not limited to lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, and commercial fish ponds).

APPLICATION RATE CHART FOR PQZ INSECTICIDE

Citrus Fruits (Crop Group 10-10)		
Australian desert lime; Australian finger lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; Mount White lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin, clementine); tangor; trifoliate orange; uniq fruit; cultivars, varieties, and/or hybrids of these		
Pest	Rate/Acre	Use Directions
Aphids	2.4 to 3.2 fl oz (0.035 to 0.047 lb ai)	<ul style="list-style-type: none">For aerial applications, apply using a minimum of 10 gallons of water per acre. USE RESTRICTIONS <ul style="list-style-type: none">Do not apply this product while bees are foraging.Do not apply this product until flowering is complete and all petals have fallen.
Citrus thrips	6.4 fl oz (0.094 lb ai)	<ul style="list-style-type: none">Do not apply through any type of irrigation system.Do not apply by Alternate Row Middle (ARM) spray method.Do not apply to citrus nurseries or citrus in greenhouses.In Florida, do not apply more than 4.8 fl oz (0.070 lb ai) per acre per year.In all other states, do not apply more than 12.8 fl oz (0.187 lb ai) per acre per year.Do not make more than 2 applications per year.Retreatment Interval: Minimum of 7 days between applicationsPreharvest Interval (PHI): 1 day

©2020 Nichino America, Inc. PQZ and Nichino America logo are registered trademarks of Nichino America, Inc.

NICHINO
AMERICA[®]

Nichino America, Inc.
4550 Linden Hill Road, Suite 501
Wilmington, DE 19808
888-740-7700