

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

PYRIFLUQUINAZON | GROUP 9B | INSECTICIDE



EPA Reg. No. 71711-37

For Use On: Whitefly in Potato

Application Method: Ground Only

State(s): Florida

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling and the EPA approved container label must be in the possession of the user at the time of application. Read the label affixed to the container for PQZ[®] Insecticide before applying. Use of PQZ Insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for PQZ Insecticide.

Potato		
Pest	Rate/Acre	Directions for Use
Whitefly	2.4 to 3.2 fl oz (0.035 to 0.047 lb ai)	Apply when pest populations are beginning to build, before reaching economic thresholds. Consult local agricultural advisor or state extension service for recommendations. <ul style="list-style-type: none">• Minimum Spray Volume: 20 gallons of water per acre• Maximum Dosage Per Acre Per Year: 4.8 fl oz (0.070 lb ai)• Maximum Applications Per Year: 2• Minimum Days Between Applications: 14 days• Preharvest Interval (PHI): 14 days• Do not feed tops to livestock.

©2019 Nichino America, Inc. PQZ is a trademark and Nichino America logo is a registered trademark of Nichino America, Inc.

NICHINO
AMERICA[®]

Nichino America, Inc.
4550 Linden Hill Road, Suite 501
Wilmington, DE 19808
888-740-7700