



PackhardTM

Guaranteed Analysis:

Calcium (Ca) 8.0%
Boron (B) 0.25%

Derived from Calcium Chloride and Boric Acid

Expiration date: 2 years after the date of manufacture.

CAUTION
KEEP OUT OF REACH OF CHILDREN

WARNING: Contains Boron (B). The use of this fertilizing material on any crops other than those listed may result in serious injury to the crops.

PRECAUTIONARY STATEMENTS:

BEFORE USING THIS PRODUCT, READ ALL PRECAUTIONS, DIRECTIONS OF USE.

CAUTION: Avoid contact with formulated product for long periods of time. Do not ingest.

IN CASE OF CONTACT WITH SKIN OR EYES: Rinse with abundant water.

GENERAL INFORMATION:

PackhardTM is a source of calcium of easy assimilation by the crop.

STORAGE AND DISPOSAL:

Store in a shaded location. Do not transfer into food or drinks containers. The remnant of product should be disposed of in accordance with local requirements.

Density: 10.85 lb/gal at 68° F

Net Content:

2.5 gal / 9.46 L

Net Weight:

27.1 lb / 12.3 kg

Lot Number:

Date of
Manufacture:



Innovak
GLOBAL

A parent company of: **Chemport inc.**

Guaranteed by: Chemport Inc.

1716 Montana Avenue El Paso TX 79902-5705

Phone: (915) 591-49-07

1. WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. THESE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER. MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, THE TYPES OF SOILS, OR THE MANNER OF USE, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER.

2. LIMITATION OF LIABILITY

IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. NEITHER MANUFACTURER NOR SELLER SHALL BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, ITS CUSTOMERS, OR USER. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT OR RETURN OF THE PRODUCT AT THE SOLE DISCRETION OF MANUFACTURER OR SELLER.

DIRECTIONS OF USE:

In general, all applications of Packhard™ must be made during the growth stage of the fruit, by spraying from fruit set to ripening, and may be continued using the specific use rate provided.

USE RATE:

CROP	DOSE	APPLICATION / TIMING
Vegetables (Peppers, tomatoes, green tomatoes, bell peppers, eggplant, etc.)	28 to 56 fl oz/ac	-2 applications of 42 fl oz/ac during the fruit development period. -42 fl oz/ac every two weeks on the harvest periods.
Cucurbits (Watermelon, melon, zucchini, cucumber, etc.)	42 to 56 fl oz/ac	-2 applications of 42 to 56 fl oz/ac during the fruit development period. -42 to 56 fl oz/ac every two weeks on the harvest periods.
Cabbage, broccoli, kale	28 to 56 fl oz/ac	-1 to 2 applications of 14 to 56 fl oz/ac during the rosette development. -2 to 3 applications of 42 to 56 fl oz/ac during the harvestable organs.
Bulb Vegetables	28 to 56 fl oz/ac	-42 fl oz/ac during the bulb formation. -1 to 2 applications of 42 to 56 fl oz/ac during the bulb growth.
Berries (Blackberry, blueberry, raspberry, strawberry, etc)	42 to 56 fl oz/ac	-2 applications of 42 fl oz/ac during the fruit development. -42 fl oz/ac every 2 weeks during the harvest period.
Ornamentals	28 to 42 fl oz/ac	-28 to 42 fl oz/ac every two weeks during the floral stem development.
Temperate fruit trees (Apples, peaches, etc.)	42 to 84 fl oz/ac	-56 to 84 fl oz/ac starting the fruit development, 2 to 3 applications after full blooming.
Grapes (Table and wine)	42 to 84 fl oz/ac	-56 to 84 fl oz/ac starting the fruit development, 4 weeks after full blooming, until veraison.
Nut crops (Almond, walnut, pistachios, etc.)	42 to 84 fl oz/ac	-56 to 84 fl oz/ac starting the fruit development, 4 weeks after full blooming, until fruit full growth.
Tropical fruit trees (Avocado, citrus, pineapple, etc.)	42 to 84 fl oz/ac	-56 to 84 fl oz/ac every two weeks starting the fruit development, four weeks after full blooming, until veraison.
Hemp	32 to 64 fl oz/ac	Foliar apply with enough water to achieve full coverage of the foliage throughout all stages of flowering at 7 to 14 day intervals between each application.

In general, all applications of Packhard™ must be made during the growth stage of the fruit, by spraying from fruit set to ripening, and may be continued using the specific use rate provided.

Information about the components of this fertilizer may be obtained by writing to Chemport Inc. and giving the lot number found on this label.

Information regarding the contents and levels of metals in this product is available on the internet at:
<http://www.aapfco.org/metals.html>

3. If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

©Chemport, Inc. All rights reserved.



Packhard™