



## SUPPLEMENTAL LABEL

### USE INSTRUCTIONS FOR SUNBURN ON ONIONS AND APPLES

#### **ONIONS**

**APPLICATION GUIDELINES:** To reduce the incidence of sunburn apply Parka at the rate of 2-4 quarts per acre. Apply in 10-50 gallons of water per acre. May be applied in a band spray. For maximum efficacy, all foliage must be covered with Parka. Do not combine with surfactants. Do not apply Parka when the air temperature is expected to meet or exceed 85 degrees Fahrenheit (29 C) during application.

**TIMING:** For best results, apply Parka preventatively. Apply prior to high heat exceeding 85 degrees Fahrenheit.

#### **APPLES**

**APPLICATION GUIDELINES:** For sunburn mitigation start applications when fruit size reaches 10-12 mm and repeat every 21 to 30 days for a total of 3 applications during the season. If treatment is for a longer season apple (i.e. Fuji or Pink Lady) or daytime temperatures remain high through harvest a 4th application is advised.

For fruit finish improvement (reduction of lenticel breakdown, flecking, scarring, russet, etc) start applications at 100% petal fall and repeat every 21 to 30 days for a total of 3 applications during the season.

To mitigate sunburn and improve fruit finish start applications at 100% petal fall and repeat every 21 to 30 days for a total of 3 to 4 applications during the season.

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE MASTER PARKA LABEL ARE TO BE FOLLOWED.**



4780 WEST HARMON AVE., SUITE 6  
LAS VEGAS, NV 89103  
TELEPHONE (888) 638-1955

**PRODUCT NAME:** PARKA™ CHERRY CUTICLE SUPPLEMENT

**MANUFACTURER:** CULTIVA LLC

**RECOMMENDED USE:** AGRICULTURE

**RESTRICTIONS ON USE:** NONE KNOWN

**EMERGENCY CONTACT / NUMBER:** (888) 638-1955 (US ONLY); (702) 987-620 (OUTSIDE US)  
PARKA IS A TRADEMARK OF CULTIVA. SURESEAL IS A TRADEMARK OF OREGON STATE UNIVERSITY.