

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Pindar™ GT

EPA Reg. No. 62719-611

### Preemergence and Postemergence Control of Annual Broadleaf Weeds in Non-Cropland

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Pindar™ GT herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Except as described in this supplemental labeling, use of Pindar GT is subject to all use precautions and limitations imposed by the label affixed to the product container.

#### Keep Out of Reach of Children

### CAUTION

#### Directions for Use

Refer to product label for Product Information, Use Precautions and Restrictions, and all other requirements not covered under this supplemental label.

#### Non-Cropland<sup>1</sup>

<sup>1</sup>Including non-food producing, non-cultivated agricultural or non-agricultural areas such as highway and utility rights-of-way, industrial sites, tank farms, storage areas, airports, fencerows not adjacent to food/feed crop fields and farmsteads

Weed Control	Rate (pints/acre)	Specific Use Directions
preemergence	3 – 4.5	Use a higher rate in the rate range for longer residual control.
postemergence		Use a lower rate in the rate range plus an approved adjuvant for control of susceptible broadleaf weeds in the early postemergence stage less than 4-leaf stage. Use a higher rate in the rate range plus an adjuvant for weeds up to 6-leaf stage. Application to weeds beyond the 6-leaf stage may result in partial control. For existing weeds not controlled by Pindar GT, a best practice is to tank mix Pindar GT with an approved postemergence herbicide for complete burndown.

#### Specific Use Precautions and Restrictions:

- Refer to Mixing Directions section on the label affixed to the product container for tank mixing precautions. Follow applicable use directions, precautions, and limitations on the respective product labels. In interpreting the labels of tank mixed products, the most restrictive label limitations must apply.
- **Preemergence:** For broader spectrum residual preemergence weed control, Pindar GT may be applied in tank mix combination with diuron, simazine or other products labeled for this use.

- **Postemergence:** For additional postemergence control of non-susceptible grass and broadleaf weeds, Pindar GT may be applied in tank mix combination with glyphosate, glufosinate or paraquat.
- Do not feed or allow animals to graze on any areas treated with Pindar GT.
- Do not apply more than a total of 4.5 pints of Pindar GT per acre in a single application.
- Pindar GT is stable on the soil surface for up to 21 days, but must be incorporated by moisture to provide effective preemergence control of susceptible weeds. A single rainfall or sprinkler irrigation of 0.5 inches or more, or flood irrigation within 21 days after application, is necessary to activate Pindar GT.

™® Trademark of Dow AgroSciences LLC

R375-002

EPA accepted 07/06/11

Replaces R375-001.