
SUPPLEMENTAL LABEL



Syngenta Crop Protection, Inc.
P. O. Box 18300
Greensboro, North Carolina 27419-8300
www.syngenta-us.com

GROUP	5	HERBICIDE
-------	---	-----------

Princep® Liquid

Herbicide

For weed control in turf and ornamental plantings

Active Ingredient:

Simazine: 2-chloro-4,6-bis(ethylamino)-s-triazine 41.9%

Other Ingredients: 58.1%

Total: 100.0%

Princep Liquid contains 4 lbs. active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-526

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

Before using Princep Liquid as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SCP 526B-S2 0508

DIRECTIONS FOR USE

TURFGRASSES FOR SOD (FL ONLY)

St. Augustinegrass, Bermudagrass, Centipedegrass, and Zoysiagrass

Apply 2-4 qts./A of Princep Liquid, according to soil texture as indicated below. This product must be watered in immediately after application if applied at a rate greater than 2 lbs. a.i./A (2 qts./A of Princep Liquid).

Muck or peat	4 qts.	Old beds	within 2 days after lifting of sod
		New beds	3-4 days after sprigging or plugging
Sandy soil	2 qts.	Old beds	within 2 days after lifting of sod
		New beds	7-10 days after sprigging or plugging

If weeds regrow, apply an additional 2 qts./A of Princep Liquid on muck or peat, or 1 qt./A of Princep Liquid on sandy soil.

Precautions: To avoid crop injury, (1) Do not apply within 30 days prior to cutting or lifting. (2) Do not apply in combination with surfactants or other spray additives. (3) Use only on turfgrass reasonably free of infestations of insects, nematodes, and diseases. (4) On newly sprigged turfgrass, temporary slowing of growth may follow application. (5) Do not exceed 4 qts. of Princep Liquid per acre (4 lbs. a.i./A) per application. (6) Do not apply more than 2 applications per year. (7) Do not exceed 6 qts. Princep Liquid per acre (6 lbs. a.i./A) per calendar year.

Note: Do not graze or feed turf clippings to animals or illegal residues may result.

TURFGRASSES FOR SOD (IN STATES EXCEPT FL)

St. Augustinegrass, Bermudagrass, Centipedegrass, and Zoysiagrass

Apply 1-2 qts./A of Princep Liquid.

Do not apply to established beds within 2 days after lifting of sod. Do not apply to new beds before 7-10 days after sprigging or plugging.

Do not apply more than 2 lbs. a.i./A (2 qts. of Princep Liquid per application) and not more than two applications per calendar year. Do not exceed 3 qts./A Princep Liquid (3 lbs. a.i./A) per calendar year.

Precautions: On newly sprigged turfgrass, hybrid bermudagrass, nondormant bermudagrass, or nondormant zoysiagrass, temporary slowing of growth and yellowing may occur following application. To avoid turf injury, (1) Use only on turfgrass reasonably free of infestations of insects, nematodes, and diseases. (2) Do not use north of NC (except in the VA Coastal Plains) or on soils with a pH above 7.8. (3) Do not use on muck or alkaline soils. (4) Do not apply over the rooting area of trees or ornamentals not listed on this label. (5) Do not seed or overseed with desirable turfgrass within 4 months before or 6 months after treatment. (6) Do not apply this product to newly seeded grasses until they have overwintered and have a well-developed rhizome system. (7) Do not exceed 2 qts./A of Princep Liquid within 12 months of seeding grasses.

Note: Do not graze or feed turf clippings to animals or illegal residues may result.

TURFGRASS FOR FAIRWAYS, LAWNS, AND SIMILAR AREAS

Bermudagrass, Centipedegrass, St. Augustinegrass, and Zoysiagrass

Apply Princep Liquid after September 1 (after October 1 for annual bluegrass) before emergence of winter annual weeds. Fall applications of Princep Liquid will control annual bluegrass, burclover, lawn burweed, common and mouseear chickweed, corn speedwell, henbit, hop clover, spurweed, and parsley-piert. Princep Liquid will also control these weeds soon after emergence. For control of summer annual weeds listed in the **General Information** section of this label, apply Princep Liquid in late winter before the weeds emerge. Apply in a minimum of 15 gals. of water per acre.

Irrigate with ½ inch of water if rainfall does not occur within 10 days after preemergence treatment.

Where annual bluegrass is the major weed, use 1 qt. of Princep Liquid per acre (22 ml. or 0.75 fl. oz. per 1,000 sq. ft.). Use 1-2 qts./A of Princep Liquid (22-44 ml. or 0.75-1.5 fl. oz. per 1,000 sq. ft.) for control of other weeds. However, do not exceed 1 qt./A of Princep Liquid per treatment on newly sprigged turfgrass or on hybrid bermudagrass.

For control of summer annuals which emerge after the initial application, apply an additional 1 qt./A of Princep Liquid at least 30 days after the initial application, but not after June 1. However, do not make more than 2 applications per calendar year.

Precautions: On newly sprigged turfgrass, hybrid bermudagrass, nondormant bermudagrass, or nondormant zoysiagrass, temporary slowing of growth and yellowing may occur following application. To avoid turf injury, (1) Use only on turfgrass reasonably free of infestations of insects, nematodes, and diseases. (2) Do not use on golf greens. (3) Do not use north of NC (except in the VA Coastal Plains) or on soils with a pH above 7.8. (4) Do not use on muck or alkaline soils. (5) Do not apply over the rooting area of trees or ornamentals not listed on this label. (6) Do not seed or overseed with desirable turfgrass within 4 months before or 6 months after treatment.

(7) Do not apply this product to newly seeded grasses until they have overwintered and have a well-developed rhizome system. (8) Do not exceed 2 qts./A of Princep Liquid within 12 months of seeding grasses. (9) Do not exceed 2 lbs. a.i./A (2 qts./A Princep Liquid) per application. (10) Do not exceed 3 qts./A Princep Liquid (3 lbs. a.i./A) per calendar year. (11) Maximum of 2 applications per year.

Note: Do not graze or feed turf clippings to animals, or illegal residues may result.

Princep® and the Syngenta logo are trademarks of a Syngenta Group Company

©2008 Syngenta

Syngenta Crop Protection, Inc.
P. O. Box 18300
Greensboro, North Carolina 27419-8300
www.syngenta-us.com