

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

## 2(ee) Recommendation



FOR DISTRIBUTION AND USE ONLY IN  
Oregon, Washington, Idaho, Montana, Utah

EPA REG. NO. 84876-2

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS  
PRODUCT IN A MANNER INCONSISTENT WITH ITS  
LABELING

This Product Bulletin Contains New or Additional  
Directions For Use Which Are Recommended By  
Sym-Agro, Inc. Under FIFRA Section 2(ee)

CROP	PEST	RATE OF APPLICATION
Pome Fruit (Crop Group 11)	Gray mold ( <i>Botrytis cinerea</i> )	40 - 45.7 fl oz/A (0.66-0.75 lb ai/A)

### DIRECTIONS FOR USE:

#### Pre-Harvest use to Control Post Harvest Disease:

Apply as a pre-harvest spray within 10 to 1 days of harvest.

Thorough coverage of the fruit is required.

Application closer to harvest may provide better efficacy.

**General:** Increase spray volume as growth increases to provide thorough coverage of foliage and fruit for optimum disease control.

### RESTRICTIONS:

Do not make more than two sequential applications of ProBlad® Verde Fungicide before alternating to a labeled fungicide with a different mode of action.

Do not make more than 5 applications of ProBlad® Verde Fungicide per harvest cycle.

### PHI: 1 Day

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED PROBLAD VERDE® FUNGICIDE (EPA REG. NO. 84876-2) LABEL MUST BE FOLLOWED. THIS RECCOMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.



Sym-Agro, Inc.  
PO Box 71  
Visalia, CA 93291

ProBlad is a trademark of CEV S.A.

ProBlad Verde\_2ee\_Pome Fruit Pre-Harvest\_07-25-2020