

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

2(ee) Recommendation:

Almonds, Stone Fruit, Grapes: Aerial Application Use Directions



IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

This Product Bulletin Contains New or Additional Directions For Use Which Are Recommended By Sym-Agro, Inc. Under FIFRA Section 2(ee)

FOR DISTRIBUTION AND USE ONLY IN
Arizona, California, Oregon, Texas, Utah, Washington

EPA REG. No. 84876-2

CROP	PEST	RATE OF APPLICATION
Grape (Crop Group 13)	Refer to the ProBlad Verde label and supplemental labels specific application guidelines and for listed diseases by crop approved in each state	18.1 to 45.7 fl oz/A (0.3 - 0.75 lb ai/A)
Almonds		
CHERRY (Crop Subgroup 12-12A), PEACH (Crop Subgroup 12-12B) and PLUM (Crop Subgroup 12-12C): Cherry subgroup: Capulin; Cherry, black; Cherry, Nanking; Cherry, sweet; Cherry, tart; cultivars, varieties, and/or hybrids of these. Peach subgroup: Peach; Nectarine; cultivars, varieties, and/or hybrids of these. Plum subgroup: Apricot; Apricot, Japanese; Jujube, Chinese; Plum; Plum, American; Plum, beach; Plum, Canada; Plum, cherry; Plum, Chickasaw; Plum, Damson; Plum, Japanese; Plum, Klamath; Plumcot; Plum, prune; Sloe; cultivars, varieties, and/or hybrids of these.		

DIRECTIONS FOR USE:

Aerial Application Use Directions: ProBlad Verde can be applied by fixed or rotary winged aircraft mix in a minimum of 15 gallons of total spray volume gallons per acre.

Optimum results are obtained when plant foliage is not too dense. COVERAGE IS ESSENTIAL FOR GOOD CONTROL. Use an output that will ensure optimum product penetration and coverage. Standard precautions should be taken to minimize spray drift. Make aerial applications when the wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Application more than 10 feet above the crop canopy is not recommended

DO NOT apply directly to water, or to areas where surface water is present

RESTRICTIONS:

Do not make more than two sequential applications of ProBlad® Verde Fungicide before alternating or tank mixing with another labeled fungicide with a different mode of action.

Do not make more than 5 applications of ProBlad® Verde Fungicide per harvest cycle.

PHI: 1 Day

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED PROBLAD VERDE® FUNGICIDE (EPA REG. No. 84876-2) LABEL MUST BE FOLLOWED. THIS RECCOMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

ProBlad is a trademark of CEV S.A.

ProBlad Verde_2ee_Aerial Application Vol. 3-24-2023