

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

2(ee) Recommendation



FOR DISTRIBUTION AND USE ONLY IN
California

EPA REG. NO. 84876-2

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS
PRODUCT IN A MANNER INCONSISTENT WITH ITS
LABELING

This Product Bulletin Contains New or Additional
Directions For Use Which Are Recommended By
Sym-Agro, Inc. Under FIFRA Section 2(ee)

| CROP | PEST | RATE OF APPLICATION |
|---------|-----------------------------------|---|
| Almonds | Hull Rot (<i>Monilinia</i> spp.) | 20.4– 45.7 fl oz/A (0.34-0.75 lb ai/A) |

DIRECTIONS FOR USE:

***Monilinia* spp.:** For control of Hull Rot apply ProBlad® Verde prior to onset of disease development. *Monilinia* Hull Rot is best managed with treatments 3-4 weeks prior to hull split. A second application should be made 7-14 days after initial hull split application to assure protection through hull split growth stage.

General: Increase spray volume as growth increases in order to provide thorough coverage of foliage and fruit for optimum disease control.

RESTRICTIONS:

Do not make more than two sequential applications of ProBlad® Verde Fungicide before alternating to a labeled fungicide with a different mode of action.

Do not make more than 5 applications of ProBlad® Verde Fungicide per harvest cycle.

PHI: 1 Day

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED PROBLAD VERDE® FUNGICIDE (EPA REG. NO. 84876-2) LABEL MUST BE FOLLOWED. THIS RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.



Sym-Agro, Inc.
PO Box 71
Visalia, CA 93291

ProBlad is a trademark of CEV S.A.

ProBlad Verde_2ee_Almond, Hull Rot_04-06-2020