

ProTransit® 50

DUAL-APPLIED LIQUID NUTRIENT



GUARANTEED ANALYSIS

Zinc(Zn)......0.5%

FOR MANUFACTURING

IISF NNIY

0.5% Chelated Zinc

Derived from: zinc EDTA.

30 US Gal / 114 Liters 8.6 lbs per Gallon @ 68° F 1.0 kgs per Liter @ 20° C Net Weight 258 lbs / 117 kgs

DO NOT STORE BELOW 40°F (5°C)

FBSciences Inc.

153 N. Main St, Suite 100 | Collierville, TN 38017 USA

V011921

F1837



BATCH/LOT NUMBER

ProTransit® 50

DIRECTIONS FOR USE

DO NOT spray in the heat of the day or when the plant is under moisture stress. DO NOT spray to the point of runoff. Use as fine of a spray mist as possible. Allow 3-4 hours after application before sprinkler irrigating to avoid washing the product off. For soil applications, use watered-in applications for best results. May be applied via irrigation system or prior to irrigation. Apply enough water to move the product into the area of active rooting, but not excessive amounts that may leach. Use the higher label rates with surface and flood irrigation. Shake well and/or agitate before use.

ProTransit 50 is compatible with most fertilizers regardless of pH. However, a standard jar test is recommended before tank mixing.

RATES & TIMING

For In Furrow or Pop-up Fertilizers mix 1.2-1.6 ounces/acre with applications near the seed.

For **Banded, Broadcast, Irrigation, & Foliar Applications** apply **1.6-2.4 ounces/acre**; higher rates of other products in the tank require the higher label rates of ProTransit 50. For foliar applications, use higher rate with spray volume over 50 gallons/acre.

For **Dry Granular Fertilizers** impregnate **6.4-48 ounces/acre** onto each ton of dry fertilizer depending on the final application of fertilizer in the field. Final applied fertilizer should carry **1.6-2.4 ounces/acre** of ProTransit 50.

WARNING

READ ENTIRE LABEL BEFORE USE | KEEP OUT OF REACH OF CHILDREN



WARNING | HAZARD STATEMENTS Harmful if swallowed. May be harmful in contact with skin.

PRECAUTIONARY STATEMENTS Wash skin thoroughly after handling. Do not eat, drink or smoke when using this product. **IF SWALLOWED:** Rinse mouth. Call a POISON CENTER or doctor/physician if you feel unwell. Dispose of contents/container in accordance with local/national regulations.