

***FIFRA Section 24(c) Special Local Need Label***  
**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON**

**Proline 480 SC Fungicide**

EPA REG No. 264-825

SLN No. WA-160004

**For Suppression of Fusarium Wilt (*Fusarium* spp.) in Spinach Grown For Seed**

**THIS LABEL FOR PROLINE 480 SC FUNGICIDE EXPIRES AND MUST NOT BE DISTRIBUTED OR USED IN ACCORDANCE WITH THIS SLN REGISTRATION AFTER DECEMBER 31, 2025.**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label. Do not apply more than 3 applications per year with a **minimum 3-week interval** between applications.

Spinach Grown For Seed Production	
PEST SUPPRESSION	RATE PER APPLICATION fluid ounces/acre
Fusarium Wilt ( <i>Fusarium</i> spp.)	5.7

**Recommended Application Timings For Spinach Seed Crops For Management Of Fusarium Wilt:**  
Apply Proline 480 SC as a banded spray or drench over the row at (or immediately after) planting. Make two repeat applications at 3-week intervals as a banded application over the rows in a large enough volume of water to drench the plants and wash the product into the soil around the roots.

**NOTES**  
Use in rotation in resistance management programs where additional products are available  
Please refer to the Proline 480 SC Fungicide Section 3 label for additional information

**RESTRICTIONS/PRECAUTIONS**

Do not apply within 14 days of harvest

Do not windrow within 14 days of last application

Maximum Proline 480 SC Fungicide allowed per year: **17.1 fl oz per acre**

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

For use under this Special Local Needs label do not apply this product through any type of irrigation system.

**Endangered Species Advisory:** This product is toxic to estuarine/marine invertebrates, and freshwater/estuaries/marine aquatic plants. Proline 480 SC Fungicide should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Natural Resources Assessment Section web site (<https://agr.wa.gov/departments/land-and-water/naturalresources/endangered-species-program>) for additional information related to pesticide use and endangered species protection. Information from EPA about no-spray buffers zones is available at <https://agr.wa.gov/departments/land-and-water/natural-resources/endangered-species-program/buffers>. Pesticide applicators may use the Washington Department of Fish and Wildlife's SalmonScape mapping tool at <https://apps.wdfw.wa.gov/salmonscape> to determine if listed salmonid species occur in or near a proposed application site.

**Crops Grown for Seed Production:**

1. All spinach seed screenings shall be disposed of in such a way that they cannot be distributed or used for human food or animal feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the director immediately upon request. Conditioner disposal records shall consist of documentation of on-farm disposal, disposal at a controlled dumpsite, incinerator, composter or other equivalent disposal site and shall include the lot numbers, amount of material disposed of, the grower(s), and the date of disposal.
2. No portion of the spinach seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.
3. Spinach seed shall bear a tag or container label which forbids use of the seed for human consumption or animal feed.
4. Spinach seed may not be distributed for human consumption or animal feed.

**WSDA Container Disposal Guidance:** Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA Waste Pesticide Program web site at <https://agr.wa.gov/departments/pesticides-and-fertilizers/pesticides/waste-pesticide>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

**IMPORTANT—READ BEFORE USE**

PROLINE 480 SC FUNGICIDE, WHEN USED ON SPINACH GROWN FOR SEED UNDER THIS SECTION 24(C) LABEL HAS NOT BEEN TESTED UNDER ALL CONDITIONS AND IN ALL TANKMIX SCENARIOS. BAYER RECOMMENDS THAT THE USER AND/OR GROWER TEST PROLINE 480 SC FUNGICIDE IN ORDER TO DETERMINE ITS COMPATIBILITY WITH THE INTENDED USE. **RISK OF CROP INJURY, CROP YIELD REDUCTION, AND/OR CROP LOSS ARE BEYOND THE CONTROL OF BAYER, AND ALL SUCH USE IS AT USER'S AND/OR GROWER'S RISK.**

This Special Conditions and Disclaimer statement is required by Bayer and is not required or enforced by the USEPA or the WSDA.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

**For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)**

*As with any crop-protection product, always read and follow label directions.*

*For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*

**Bayer CropScience LP  
800 N.Lindbergh Blvd.  
St. Louis, MO 63167  
1-866-99BAYER (1-866-992-2937)**