

# Supplemental Labeling



Dow AgroSciences LLC      9330 Zionsville Road      Indianapolis, IN 46268-1054 USA

## PropiMax<sup>®</sup> EC EPA Reg. No. 62719-346

This supplemental label expires June 30, 2013 must not be used or distributed after this date.

### Disease Control on Bushberries (Subgroup 13-07B)<sup>1</sup>, Caneberries (Subgroup 13-07A)<sup>2</sup>, and Low Growing Berries (Subgroup 13-07G) (Except Cranberry)<sup>3</sup>

**ATTENTION**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for PropiMax<sup>®</sup> EC fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of PropiMax EC according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for PropiMax EC.

### Directions for Use

<sup>1</sup>Bushberries (subgroup 13-07B) including aronia berry, black currant, buffalo currant, Chilean guava, edible honeysuckle, elderberry, European barberry, gooseberry, highbush blueberry, highbush cranberry, huckleberry, jostaberry, juneberry, lingonberry, lowbush blueberry, native currant, red currant, salal, sea buckthorn, cultivars, varieties, and/or hybrids of these

<sup>2</sup>Caneberries (subgroup 13-07A) including bingleberry, blackberry, black raspberry, boysenberry, dewberry, loganberry, lowberry, marionberry, olallieberry, red raspberry, wild raspberry, youngberry, cultivars, varieties and/or hybrids of these

<sup>3</sup>Low growing berries (subgroup 13-07G) except cranberry) including bearberry, bilberry, lowbush blueberry, cloudberry, lingonberry, muntries, partridgeberry, strawberry, cultivars, varieties, and/or hybrids of these

Refer to product label for Integrated Pest Management, Fungicide Resistance Management, Spray Equipment, Mixing Directions, and Application Directions.

#### Bushberries (Subgroup 13-07B) and Caneberries (Subgroup 13-07A)

Apply PropiMax EC by either ground equipment in a minimum of 15 gpa or aerial equipment in a minimum of 5 gpa.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
leaf spot, stem canker	<i>Septoria albopunctata</i>	6	Apply when conditions favor disease development. Reapply on a 4-week interval.	<ul style="list-style-type: none"> <li>• Do not apply more than a total of 30 fl oz of PropiMax EC (0.84 lb active ingredient) per acre per season.</li> <li>• <b>Preharvest Interval:</b> Do not apply within 30 days of harvest.</li> </ul>
rust	<i>Pucciniastrum vaccinii</i>		Make the first application any time prior to bloom and again after petal fall. If needed, reapply just after harvest.	
leaf spot	<i>Septoria</i> spp.		Apply as a delayed dormant spray after	
leaf and cane spot	<i>Septoria rubi</i>			

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
			training in the spring. Reapply in late spring, at bud break, and once flowering has begun.	
mummyberry disease	<i>Monilinia vaccinicorymbosi</i>		Make first application beginning at green tip and repeat in 7 to 10 days. If conditions are favorable for disease development, reapply at pink bud and every 7 to 10 days through petal fall.	
powdery mildew	<i>Microsphaera vaccinii</i>		Apply at 5 to 10% bloom. Reapply at full bloom and on a 14-day interval while conditions favor disease development.	

**Low Growing Berries (Subgroup 13-07G) (Except Cranberry)**

Apply PropiMax EC by either ground equipment in a minimum of 20 gpa or aerial equipment in a minimum of 15 gpa.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
anthracnose	<i>Colletotrichum acutatum</i>	4	Apply when disease levels are no more than 5%. Apply up to 4 times on a 7-day interval. Do not make more than 2 consecutive applications of this product before rotating to another fungicide with a different mode of action.	<ul style="list-style-type: none"> <li>Do not apply more than a total of 16 fl oz of PropiMax EC (0.45 lb active ingredient) per acre per season.</li> <li><b>Preharvest Interval:</b> There is no preharvest interval.</li> </ul>
leaf rust	<i>Phragmidium potentillae</i>			
leaf spot	<i>Cercospora fragariae</i>			
powdery mildew	<i>Sphaerotheca maculeris</i>			

**Note:** This pesticide is toxic to fish. Refer to the Environmental Hazards section of the label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R142-002

EPA accepted 01/05/12

Initial printing