

FIFRA §2(ee) Recommendation

Propicure 3.6F

Fungicide
EPA Reg. No. 83222-18

For The Control of Rust and Scab on Almonds in California Only

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- The user must refer to the Propicure 3.6F Fungicide container label and read and follow all directions for use, restrictions, and precautions found on that label.
- Users should have this recommendation in their possession at the time of use.
- This recommendation is permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the US EPA.

CROP: ALMONDS

| PESTS CONTROLLED | RATE OF APPLICATION PER ACRE | APPLICATION INSTRUCTIONS |
|--|-------------------------------------|---|
| Rust (<i>Tranzschelia discolor</i>) Scab (<i>Cladosporium carpophilum</i>) | 6 to 8 fl. oz. | Apply 6 to 8 fl. oz (0.18- 0.225 lbs active) per acre. Begin application approximately 2-5 weeks after petal fall when environmental conditions are favorable for disease development. For optimum disease control, apply on a 7- to 14-day schedule. For disease management, make no more than two consecutive applications before rotation to a fungicide with a different mode of action (not a Group 3) |
| <ul style="list-style-type: none">• Do not apply Propicure 3.6F within 60 days of harvest.• Do not apply more than 32 fl. oz. (0.90 lb. ai) of Propicure 3.6F per acre per season.• Do not apply more than 0.90 lbs active of any propiconazole product per acre per season.• Do not graze livestock in treated areas or cut treated cover crop for feed. | | |

Effective Date: 1-7-13

Direct Ag Source, LLC
30473 260th Street, Eldora, IA 50627