



## Section 2(ee) Recommendation

**Date:** January 5, 2011

**Product:** Quadris Top™  
EPA Registration No. 100- 1313

**Uses:** Addition of sweet orange scab (*Elsinoe australis*) as a disease controlled in citrus

**States:** Florida, Louisiana

### Directions for Use

Crop	Target Disease	Use Rate Fl. oz. Product/A	Remarks
<b>Citrus</b>  Grapefruit Lemon Lime Orange (sour and sweet) Tangerine  Including all cultivars and/or hybrids of these	<b>Sweet Orange            Scab            (<i>Elsinoe australis</i>)</b>	10 - 15.4	Quadris Top applications should begin prior to disease development and continue throughout the season on 7-21 day intervals following the resistance management guidelines. Applications may be made by ground or air. An adjuvant may be added at recommended rates. A horticultural spray oil should be used to improve control of greasy spot.
			If disease pressure is high, use the shortest interval and highest rate.
<b>Application:</b> For best results, sufficient water volume must be used to provide thorough coverage. Quadris Top can be applied by either ground or aerial application. A minimum of 15 gals./A for ground applications is recommended. For aerial applications a minimum of 10 gals./A of water is recommended.			Make no more than 2 sequential applications before alternating to another fungicide with a non-QoI (Group 11) different mode of action. Do not make more than 4 applications of Quadris Top or other Group 11 fungicides per season.

**Complete list of citrus fruits:** Calamondin; citrus citron; citrus hybrids (includes chironja, tangelo, tangor); grapefruit; kumquat; lemon; lime; mandarin (tangerine); orange, sour; orange, sweet; pummelo; Satsuma mandarin

#### Specific Use Restrictions:

- Do not use Quadris Top in citrus plant propagation nurseries.
- Do not apply more than 61.5 fl. ozs./A of Quadris Top per season.
- Do not apply more than 0.5 lb. ai/A per crop of difenoconazole containing products.
- Do not apply more than 1.5 lbs. ai/A per crop of azoxystrobin containing products.
- May be applied the day of harvest (0 day PHI).

**The user must refer to the federally approved Quadris Top label and read and follow all directions for use, restrictions, and precautions. It is a violation of federal law to use this product in a manner inconsistent with its labeling. The user should have this recommendation in their possession at the time of use.**

**This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the EPA.**

Syngenta Crop Protection, Inc.  
Greensboro, NC 27419

Quadris Top™ - Trademark of a Syngenta group company

FL1313051AC0111