

---

## SUPPLEMENTAL LABELING

---

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300  
**SCP 1178A-S16 0614**

**GROUP 3 11 FUNGICIDES**

### **Quilt® Fungicide**

Broad spectrum fungicide for control of plant diseases

**This supplemental label expires on June 13, 2017 and must not be used or distributed after this date.**

Active Ingredient:

Azoxystrobin.....	7.0%
Propiconazole .....	11.7%
<hr/>	
Other Ingredients:	81.3%
Total:	100.0%

Quilt Fungicide is a suspoemulsion (SE) formulation.

Quilt Fungicide contains 1.04 lb ai propiconazole and 0.62 lb ai azoxystrobin per gallon.

**KEEP OUT OF REACH OF CHILDREN.**

## **WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**EPA Reg. 100-1178**

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Quilt Fungicide as permitted according to this Supplemental Labeling, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**

**syngenta**

## DIRECTIONS FOR USE

**Rotational Crops:** Soybeans may be planted as a double crop following a cereal crop which has been treated with Quilt. Do not use hay, forage, or fodder from the soybean crop as any component of animal feed or bedding. Plant wild rice 45 days after Quilt application. To avoid possible illegal residues, do not plant any other crop intended for food, grazing, or any component of animal feed or bedding within 105 days of Quilt application to the preceding crop, unless the second crop appears on this label.

<b>Rotational Crops</b>	<b>Planting Time From Last Quilt Application</b>
Bean, dry and succulent Bulb crops Berries, bushberry and caneberry subgroups Carrots Celery (and other leaf petiole crops - subgroup 4B) Cereals (wheat, barley, triticale, oat and rye) Corn (field, seed, popcorn, and sweet) Grasses grown for seed Mint Peanuts Rice Sorghum Soybeans Strawberries Sugar beets	0 days
Buckwheat Millet	12 Months
Wild rice	45 days
Alfalfa (if propiconazole rate does not exceed 0.22 lb ai/acre/season)	75 days
All Other Crops Intended for Food and Feed	105 days

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
<b>Corn</b> Field Pop (Includes Seed Production)	Northern Corn Leaf Blight ( <i>Setosphaeria            turcica</i> ) Northern Corn Leaf Spot ( <i>Cochliobolus            carbonum</i> ) Southern Corn Leaf Blight ( <i>Cochliobolus            heterostrophus</i> )  also known as Helminthosporium Leaf Blights ( <i>H. maydis</i> , <i>H. turcicum</i> , <i>H. carbonum</i> )	7 - 14	For leaf blights apply Quilt when disease first appears. Continue on a 7- to 14-day schedule. Use the low rate when disease pressure is low. Under heavy disease pressure or if conditions are favorable for disease, apply the highest labeled rate.  Apply no more than 2 applications of Quilt or any other Group 11 fungicide per year.  Do not use adjuvants or other additives after the V8 growth stage and prior to the VT growth stage, as use during these development times may impose stress on the plant that could inhibit proper kernel development. VT is defined as when the last branch of the tassel is completely visible but silks have not yet emerged from the ear shoot.
	Anthracnose Leaf Blight ( <i>Colletotrichum            graminicola</i> ) Eye Spot ( <i>Aureobasidium            zeae</i> ) Gray Leaf Spot ( <i>Cercospora zeae-            maydis</i> ) Rusts ( <i>Puccinia spp.</i> )	10.5 - 14	Apply Quilt when disease first appears. If conditions favorable for disease persist, continue to apply on a 14-day schedule. Under heavy disease pressure or if conditions are favorable for disease, apply the highest labeled rate.  Apply no more than 2 applications of Quilt or any other Group 11 fungicide per year.

**Application:** Quilt may be applied by ground, air or chemigation. For best results, sufficient coverage is very important. Use of a crop oil concentrate is recommended for aerial applications to reduce evaporation and enhance canopy penetration and coverage. Consult your aerial applicator for recommended concentration of crop oil concentrate. Use higher water volumes for aerial applications if equipment and/or conditions will not provide good coverage.

**Specific Use Restrictions:**

- 1) Do not apply more than 56 fl oz/A/season of Quilt.
- 2) Do not apply more than 28 fl oz (0.224 lb ai propiconazole) for field corn harvested for forage.
- 3) Do not apply more than 0.45 lb ai propiconazole-containing products/A/season.
- 4) Do not apply more than 2.0 lb ai azoxystrobin-containing products/A/season.
- 5) Do not apply within 30 days of harvest (30-day PHI) for forage, grain, or stover.

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
<b>Corn, Sweet</b> Sweet corn (Includes Seed Production)	Northern Corn Leaf Blight ( <i>Setosphaeria turcica</i> ) Northern Corn Leaf Spot ( <i>Cochliobolus carbonum</i> ) Southern Corn Leaf Blight ( <i>Cochliobolus heterostrophus</i> )  also known as Helminthosporium Leaf Blights ( <i>H. maydis</i> , <i>H. turcicum</i> , <i>H. carbonum</i> )	7 - 14	For leaf blights apply Quilt when disease first appears. Continue on a 7- to 14-day schedule. Use the low rate when disease pressure is low. Under heavy disease pressure or if conditions are favorable for disease, apply the high rate.  Alternate applications of Quilt with Tilt or another product with a different mode of action than Group 11 fungicides.
	Anthracnose Leaf Blight ( <i>Colletotrichum graminicola</i> ) Eye Spot ( <i>Aureobasidium zeae</i> ) Gray Leafspot ( <i>Cercospora zeae- maydis</i> ) Rusts ( <i>Puccinia spp.</i> )	10.5 - 14	Apply Quilt when disease first appears. If conditions favorable for disease persist, continue to apply on a 14-day schedule.  Alternate applications of Quilt with Tilt or another product with a different mode of action than Group 11 fungicides.

**Application:** Quilt may be applied by ground, air or chemigation. For best results, sufficient coverage is very important. Use of a crop oil concentrate is recommended for aerial applications to reduce evaporation and enhance canopy penetration and coverage. Consult your aerial applicator for recommended concentration of crop oil concentrate. Use higher water volumes for aerial applications if equipment and/or conditions will not provide good coverage.

**Specific Use Restrictions:**

- 1) Do not apply more than 56 fl oz/A/season of Quilt.
- 2) Do not apply more than 0.45 lb ai propiconazole-containing products/A/season.
- 3) Do not apply more than 2.0 lb ai azoxystrobin-containing products/A/season.
- 4) Do not apply to sweet corn within 14 days of harvest (14-day PHI) for ears or forage.

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
Soybeans	Aerial Web Blight ( <i>Rhizoctonia solani</i> ) Alternaria Leafspot ( <i>Alternaria</i> spp.) Anthracnose ( <i>Colletotrichum truncatum</i> ) Brown Spot ( <i>Septoria glycines</i> ) Cercospora Blight and Leaf Spot ( <i>C. kickuchii</i> ) Frogeye Leaf Spot ( <i>Cercospora sojina</i> ) Pod and Stem Blight ( <i>Diaporthe</i> spp.) Soybean Rust ( <i>Phakopsora pachyrhizi</i> )	14 – 20.5	<p>Aerial Web Blight: Apply 14- 20.5 fl oz/A at the first appearance of disease and repeat the application 14-21 days later. Under severe disease conditions use the higher rate and shorter interval.</p> <p>Other foliar diseases (except rust): Apply 20.5 fl oz/A at growth stage R3 (early pod set when pods are 1/8-1/4 inch long) and 14-21 days later at growth stage R5 (pod fill).</p> <p>Soybean Rust: Apply 14-20.5 fl oz/A at first indication that disease is in the area. For best control, preventive applications work best. Repeat on a 14- to 21-day interval. Use higher rate and shorter interval when disease is present in the field and incidence is less than 2% (2 plants in 100 are infected). If incidence is greater than this or if disease is in mid canopy, control will not be acceptable. Scouting for the disease and/or being aware of the proximity of the disease via monitoring systems will aid in the proper timing to maximize the effectiveness of the fungicide applications.</p> <p>On certain varieties, Quilt applications may cause crinkled, smaller and/or greener leaves. Yields of beans displaying these characteristics have not been reduced due to Quilt treatments.</p>

**Application:** Quilt is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions will not provide good coverage. Quilt may be applied by ground, air or chemigation.

**Specific Use Restrictions:**

- 1) Do not apply more than 42 fl oz/A/season of Quilt.
- 2) Do not apply more than 0.34 lb ai of propiconazole-containing products/A/season.
- 3) Do not apply more than 1.5 lb ai of azoxystrobin-containing products/A/season.
- 4) Apply up to Stage R6.

Quilt® and the Syngenta logo are trademarks of a Syngenta Group Company.

©2014 Syngenta