
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 1178A-S17 0216

GROUP	3 11	FUNGICIDES
-------	-------------	------------

Quilt® Fungicide

Broad spectrum fungicide for control of plant diseases

This supplemental label expires on February 1, 2019 and must not be used or distributed after this date.

Active Ingredient:

Azoxystrobin.....	7.0%
Propiconazole	11.7%
<hr/>	
Other Ingredients:	81.3%
Total:	100.0%

Contains 1.04 lb ai propiconazole and 0.62 lb ai azoxystrobin per gallon.

Quilt Fungicide is formulated as a suspo-emulsion (SE).

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

EPA Reg. 100-1178

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Quilt Fungicide as permitted according to this Supplemental Labeling, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

Rotational Crops	Planting Time From Last Quilt Application
Brassica, leafy greens (subgroup 5B) Bulb Crops Carrots Celery (and other leaf petiole crops - subgroup 4B) Cereals (wheat, barley, triticale) Corn (field, seed, popcorn, and sweet) Dill Grasses Grown for Seed Mint Oats Peanuts Quinoa Radish Rice Rye Sorghum Soybeans Strawberries Sugar Beets Watercress	0 days
Buckwheat Millet	12 Months
Wild rice	45 days
Alfalfa (if propiconazole rate does not exceed 0.22 lb ai/acre/season)	75 days
All Other Crops Intended for Food and Feed	105 days

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
Brassica, leafy greens (subgroup 5B) Cabbage, Chinese (bok choy) Collards Kale Mustard Greens (See additional crops below)	Alternaria Leaf Spot <i>(Alternaria spp.)</i> Anthracnose <i>(Colletotrichum spp.)</i> Black Spot <i>(Alternaria spp.)</i> Cercospora Leaf Spot <i>(Cercospora spp.)</i> Powdery Mildew <i>(Erysiphe polygoni)</i> Ring Spot <i>(Mycosphaella brassicicola)</i>	10.5 - 14	Begin applications at first sign of disease. Repeat on a 7- to 10-day interval. Apply no more than two applications before switching to another fungicide with a different mode of action. If disease levels continue to increase, immediately switch to a fungicide with a different mode of action.

Leafy Brassica (subgroup 5B): Broccoli raab; Cabbage, Chinese (bok choy); Collards, Kale, Mizuna, Mustard greens; Mustard spinach; Rape greens

Application: Quilt is most effective when applied and allowed to dry before a rainfall. For best results, sufficient water volume should be used to provide thorough coverage. Quilt may be applied by ground, air, or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 42 fl oz/A/season of Quilt.
- 2) Do not apply more than 0.34 lb ai propiconazole-containing product/A/season.
- 3) Do not apply more than 0.75 lb ai azoxystrobin-containing product/A/season.
- 4) Do not apply within 7 days of harvest (7-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
Dill	Cercospora Leaf Blight <i>(Cercosporidium punctum)</i> Powdery Mildew <i>(Erysiphe heraclei)</i>	10.5 - 14	Begin applications at first sign of disease. Repeat on a 7- to 10-day interval. If disease levels continue to increase, immediately switch to a fungicide with a different mode of action.

Application: Quilt is most effective when applied and allowed to dry before a rainfall. For best results, sufficient water volume should be used to provide thorough coverage. Quilt may be applied by ground, air, or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 42 fl oz/A/season of Quilt.
- 2) Do not apply more than 0.34 lb ai propiconazole-containing product/A/season.
- 3) Do not apply more than 1.5 lb ai azoxystrobin-containing product/A/season.
- 4) Do not apply within 7 days of harvest (7 day PHI).

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
Quinoa	Leaf Spot (<i>Ascochyta hyalospora</i>) Stalk Rot (<i>Phoma exigua</i>)	10.5 – 14	Apply prior to disease development. An adjuvant may be added at recommended rates.

Application Directions: Quilt can be applied by ground, chemigation, or aerial application.

Specific Use Restrictions:

- 1) Do not apply more than 2 applications/A/season.
- 2) Do not apply more than 28.0 fl oz/A/season of Quilt.
- 3) Do not apply more than 0.22 lb ai propiconazole-containing products/A/season.
- 4) Do not apply more than 0.40 lb ai azoxystrobin-containing products/A/season.
- 5) Under certain environmental conditions, tank mixes of Quilt plus herbicides and/or fertilizers may cause crop injury.
- 6) Do not apply within 7 days of harvest (7-day PHI) for forage and hay.
- 7) Do not apply within 30 days of harvest (30-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
Radish	Cercospora Leaf Spot (<i>Cercospora</i> spp.)	10.5 - 14	Begin applications at first sign of disease. Repeat on a 7- to 10-day interval. If disease levels continue to increase, immediately switch to a fungicide with a different mode of action.

Application: Quilt is most effective when applied and allowed to dry before a rainfall. For best results, sufficient water volume should be used to provide thorough coverage. Quilt may be applied by ground, air, or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 56 fl oz/A/season of Quilt.
- 2) Do not apply more than 0.45 lb ai propiconazole-containing product/A/season.
- 3) Do not apply more than 2.0 lb ai azoxystrobin-containing product/A/season.
- 4) Do not apply within 14 days of harvest (14 day PHI).

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
Watercress	Alternaria Leaf Spot (<i>Alternaria</i> spp.) Cercospora Leaf Spot (<i>C. nasturtii</i>)	10.5 - 14	Begin applications at first sign of disease. Repeat on a 7-10 day interval. Make no more than 2 applications before harvesting leaves. Up to 4 applications can be made per season. If disease levels continue to increase, immediately switch to a fungicide with a different mode of action.

Application: Quilt is most effective when applied and allowed to dry before a rainfall. For best results, sufficient water volume should be used to provide thorough coverage. Quilt may be applied by ground, air, or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 56 fl oz/A/season of Quilt.
- 2) Do not apply more than 0.45 lb ai propiconazole-containing product/A/season.
- 3) Do not apply more than 1.5 lb ai azoxystrobin-containing product/A/season.
- 4) Do not apply within 7 days of harvest (7 day PHI).

Quilt Rate Conversion Table

Fl oz/A	Lb ai/A Propiconazole	Lb ai/A Azoxystrobin	Treated acres per gallon product
3	0.025	0.015	42.6
4	0.033	0.020	32.0
7	0.056	0.034	18.3
10.5	0.085	0.051	12.2
14	0.113	0.070	9.1
15	0.122	0.073	8.5
20.5	0.167	0.100	6.2
27.5	0.224	0.134	4.6
34.5	0.280	0.169	3.7

Quilt® and the Syngenta logo are trademarks of a Syngenta Group Company.

©2016 Syngenta