

QUINOXYFEN

GROUP

13

FUNGICIDE

SUPPLEMENTAL LABELING

This supplemental label expires October 14, 2024 and must not be used or distributed after this date.



EPA Reg. No. 33906-26-10163

ACTIVE INGREDIENT: Quinoxifen	% By Wt.
5,7-dichloro-4-(p-fluorophenoxy) quinoline	22.58%
OTHER INGREDIENTS**	77.42%
TOTAL	100.00%

Contains 2.08 lbs. of active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION/ PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label must be followed.
- This labeling must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for Quintec Fungicide before applying.

APPLICATION DIRECTIONS

PREHARVEST INTERVAL

The required days between the last application and harvest are given in () after each crop name

CROP	DISEASE	RATE (Fl. Oz./acre)	INSTRUCTIONS
CHERRY (7)	Powdery mildew (<i>Sphaerotheca pannosa</i> , <i>Podosphaera leucotricha</i>)	8.7	Apply Quintec before visible symptoms of powdery mildew appear. Quintec will not control latent or established infections of powdery mildew. If powdery mildew infection is established, apply Quintec in tank mix combination with a curative fungicide or following application of a curative spray or, if multiple diseases are present, following application of a broad-spectrum fungicide. Under low disease conditions (as per a predictive modeling system), minimum label use rates per application can be used. For optimum results, use maximum label rates and short application intervals for severe or threatening disease conditions. Ground: Apply uniformly by ground application in a minimum spray volume of 30 gallons per acre. Disease control may be reduced at low spray volumes or if spray coverage is inadequate. DO NOT apply under conditions when uniform spray coverage cannot be obtained or when spray drift may occur. Aerial: DO NOT aerially apply this product on cherry, grape, hops, and stone fruits.
Restrictions for Cherry: <ul style="list-style-type: none"> • Preharvest Interval: DO NOT apply within 7 days of harvest. • Maximum Application Rate: DO NOT apply more than 17.4 fl. oz. (0.283 lbs./ai/acre) of Quintec per acre per calendar year. • Maximum Single Application Rate: 8.7 fl. oz. / acre (0.141 lbs./ai/acre). • Minimum Re-treatment Interval: 10 days. • Maximum Applications/year: 2 • Quintec is in mode of action Group 13. DO NOT make more than two applications of Quintec per calendar year. 			

Gowan[®]
The Go To Company

Distributed by:
Gowan Company, LLC
P.O. Box 5569
Yuma, AZ. 85366-5569