

**FIFRA Section 24(c) Special Local Need Label
FOR DISTRIBUTION AND USE ONLY WITHIN THE
STATE OF OREGON**

ROMEO

EPA Reg. No. 91810-2
EPA SLN No. OR-210011

**For Control of Downy Mildew and
Powdery Mildew in Hops**

This label is valid until December 31, 2022 or until otherwise amended, withdrawn, canceled, or suspended.

Active Ingredient:

Cerevisane (cell walls of *Saccharomyces cerevisiae* strain LAS117) 94.1%

Other Ingredients 5.9%

Total 100.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

ENVIRONMENTAL HAZARDS

For terrestrial uses. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This FIFRA Special Local Need (SLN) label and the ROMEO EPA-registered label for this product must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the main EPA-registered label and this SLN label.
- Refer to the main EPA-registered label concerning early re-entry into treated areas.

PRODUCT INFORMATION

ROMEO is a Systemic Resistance Inducer (SRI) that acts preventively. Its active ingredient strongly induces plant defense mechanisms, and so prepares the plant to defend itself against fungal and bacterial attacks.

APPLICATION INSTRUCTIONS

- Apply ROMEO as a foliar spray using conventional spray equipment.
- The recommended range of spray solution is 20-100 gallons per acre.
- Apply to the point of saturation, using a sufficient volume of spray solution to ensure complete coverage of plant foliage without run-off.
- When possible, apply the product on both the top and underside of the leaves.
- Apply at a spray interval of 7-10 days depending on disease pressure.

APPLICATION RATE

CROP	TARGET DISEASES	APPLICATION RATE
HOPS	Downy Mildew, Powdery Mildew	0.45-0.68 lbs./acre

RESTRICTIONS AND PRECAUTIONS

- If possible, apply the product early in the morning; avoid application during periods of hot temperatures.
- For use under this SLN, do not apply through any type of chemigation.
- If heavy rain occurs within 48 hours after treatment, repeat the application.
- In the case of high disease pressure/risk, ROMEO should be used in a program with other fungicides.
- Pre-harvest interval (PHI) = 0 days.

2021_0617

Section 24(c) Registrant



Lesaffre Yeast Corporation
7475 W. Main Street
Milwaukee, WI 53214