

**FIFRA Section 24(c) Special Local Need Label**  
**FOR DISTRIBUTION AND USE ONLY WITHIN THE**  
**STATE OF WASHINGTON**

**ROMEO**

EPA Reg. No. 91810-2  
EPA SLN No. WA-210007

**For Control of Downy Mildew and**  
**Powdery Mildew in Hops**

This label for Romeo expires and must not be distributed or used in accordance with this SLN registration after December 31, 2022.

**Active Ingredient:**

Cerevisane (cell walls of <i>Saccaromyces cerevisiae</i> strain LAS117) .....	94.1%
Other Ingredients .....	5.9%
Total .....	100.0%

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

**ENVIRONMENTAL HAZARDS**

For terrestrial uses. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

### **DIRECTIONS FOR USE**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This FIFRA Special Local Need (SLN) label and the ROMEO EPA-registered label for this product must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the main EPA-registered label and this SLN label.
- Refer to the main EPA-registered label concerning early re-entry into treated areas.

### **PRODUCT INFORMATION**

ROMEO is a Systemic Resistance Inducer (SRI) that acts preventively. Its active ingredient strongly induces plant defense mechanisms, and so prepares the plant to defend itself against fungal and bacterial attacks.

### **APPLICATION INSTRUCTIONS**

- Apply ROMEO as a foliar spray using conventional spray equipment.
- The recommended range of spray solution is 20-100 gallons per acre.
- Apply to the point of saturation, using a sufficient volume of spray solution to ensure complete coverage of plant foliage without run-off.
- When possible, apply the product on both the top and underside of the leaves.
- Apply at a spray interval of 7-10 days depending on disease pressure.

### **APPLICATION RATE**

<b>CROP</b>	<b>TARGET DISEASES</b>	<b>APPLICATION RATE</b>
<b>HOPS</b>	Downy Mildew, Powdery Mildew	0.45-0.68 lbs./acre

### **RESTRICTIONS AND PRECAUTIONS**

- If possible, apply the product early in the morning; avoid application during periods of hot temperatures.
- For use under this SLN, do not apply through any type of chemigation.
- If heavy rain occurs within 48 hours after treatment, repeat the application.
- In the case of high disease pressure/risk, ROMEO should be used in a program with other fungicides.
- Pre-harvest interval (PHI) = 0 days.

2021-0809

Section 24(c) Registrant

Lesaffre Yeast Corporation  
7475 W. Main Street  
Milwaukee, WI 53214