

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Rally® 40W

EPA Reg. No. 62719-411

EPA 24(c) Special Local Need Registration SLN OR-060011
For Distribution and Use Only in the State of Oregon

Expiration Date: This label valid until December 31, 2008 or until otherwise amended, withdrawn, canceled, or suspended.

Blackberry Rust Control in Evergreen Blackberry (Thornless Evergreen)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label and the federal label must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for Rally® 40W fungicide before applying. Follow all other applicable directions, restrictions, and precautions on the EPA-registered label.
- Use of Rally 40W according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Rally 40W.

Active Ingredient:

myclobutanil: a-butyl-a-(4-chlorophenyl)-1H-1,2,4-triazole-1-propanenitrile	40%
Inert Ingredients	60%
Total	100%

WARNING

AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Environmental Hazards

For terrestrial uses, do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply when weather conditions favor drift or runoff from areas treated.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant gloves (Category A in EPA's chemical-Resistant Category Selection Chart)
- Shoes plus socks
- Protective eyewear

General Information

Rally 40W is a systemic, protectant and curative fungicide. Optimum disease control is achieved when the fungicide is applied in a regularly scheduled preventative spray program.

Blackberry rust (*Phragmidium violaceum*) spores are carried by the wind. When *P. violaceum* infects leaves, purple spots develop on the top of leaves with corresponding yellow and black pustules underneath the leaves. Other plant tissues can become infected, including floral bracts. Some defoliation and die-back of the cane can also occur. As part of a comprehensive IPM program, susceptible clones of Himalayan blackberry plants, which may serve as a source of inoculum, should be controlled.

Application Directions

Application Timing: Initiate applications when fully expanded leaves are present. Consult your Dow AgroSciences representative, extension specialist, or certified crop advisor for any additional local use recommendations for your area.

Application Rate: Apply as a foliar spray at the rate indicated for the target disease.

Disease	Rally 40W (oz/acre)	Use Recommendations	Restrictions
blackberry rust (<i>Phragmidium violaceum</i>)	5 (2 oz ai/acre)	Apply at 10- to 18-day intervals. Use the shorter spray interval under heavy disease pressure.	<ul style="list-style-type: none">• Do not apply more than 20 oz of Rally 40W (0.5 lb active) per acre per calendar year.• Do not make more than 4 applications per calendar year. One application may be made after harvest.• Pre-harvest Interval: Do not apply within 5 days of harvest.

®Trademark of Dow AgroSciences LLC

R867-042

Accepted: 03/22/06

Initial printing.