



PEEL FILM HERE ↑

RAMPART 6.8

A fungicide for the control of phytophthora, downey mildew, pythium, and other labeled diseases on agricultural and greenhouse crops, indoor and outdoor ornamentals, bedding plants, turf and commercial forestry and domestic trees including conifers, pines, oaks, and post-harvest agricultural treatment.

ACTIVE INGREDIENT:

Mono- and dipotassium phosphite* 57.0%

OTHER INGREDIENTS: 43.0%

TOTAL **100.0%**

*Rampart 6.8 contains 6.78 lbs./gallon of the active ingredients, mono- and di-potassium phosphite, equivalent to 4.16 pounds phosphorous acid per gallon (35% by weight).

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center (NPIC) at 1-800-858-7378 for emergency medical treatment information.

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

See inside label booklet for additional Precautionary Statements, and Directions for Use including Storage and Disposal instructions.

ENVIRONMENTAL HAZARDS

For terrestrial uses. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

EPA REG. NO. 82940-1-34704

061820 V1D 06W20

FORMULATED FOR
LOVELAND PRODUCTS, INC.® P.O. BOX 1286 GREELEY, COLORADO 80632-1286



RAMPART 6.8

A fungicide for the control of phytophthora, downey mildew, pythium, and other labeled diseases on agricultural and greenhouse crops, indoor and outdoor ornamentals, bedding plants, turf and commercial forestry and domestic trees including conifers, pines, oaks, and post-harvest agricultural treatment.

ACTIVE INGREDIENT:

Mono- and di-potassium phosphite* 57.0%

OTHER INGREDIENTS: 43.0%

TOTAL 100.0%

*Rampart 6.8 contains 6.78 lbs./gallon of the active ingredients, mono- and di-potassium phosphite, equivalent to 4.16 pounds phosphorous acid per gallon (35% by weight).

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center (NPIC) at 1-800-858-7378 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

See inside label booklet for additional Precautionary Statements, and Directions for Use including Storage and Disposal instructions.

ENVIRONMENTAL HAZARDS

For terrestrial uses. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

EPA REG. NO. 82940-1-34704

061820 V1D 06W20

FORMULATED FOR

LOVELAND PRODUCTS, INC.®, P.O. BOX 1286, GREELEY, COLORADO 80632-1286

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides 40 CFR 170.240(d) (4-5), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using this product. Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water), is:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

GENERAL APPLICATION INSTRUCTIONS

Rampart 6.8 may be applied as a foliar spray, trunk injection, soil drench, soil incorporation, or bare root dip. For foliar sprays, apply Rampart 6.8 with sufficient water for adequate coverage of foliage, according to crop and growth stage. To ensure good coverage, spray until wet but not to the point of runoff.

LOW VOLUME AND ULTRA-LOW VOLUME APPLICATIONS

Label rates are written for conventional application equipment. For LV/ULV equipment, use the per acre rate of Rampart 6.8 and adjust amount of water to equipment needs. NOTE: Always conduct a phytotoxicity test on a small area before applying to a large area in order to assess any potential risk to plants.

MIXING INSTRUCTIONS

1. Fill the spray tank with ½ of the required volume of water.
2. Begin agitation (either hydraulic or mechanical) and add Rampart 6.8 slowly to the tank.
3. Continue agitation and fill the tank to the desired volume with water.
4. Maintain agitation while applying.

COMPATIBILITY

Crop injury may result from mixing Rampart 6.8 with certain surfactants, foliar fertilizers or other pesticides. Because Rampart 6.8 is a slightly acidic buffer solution users should avoid mixing Rampart 6.8 with strongly acidic or alkaline materials. Do not tank mix without first testing the mixture's compatibility nor apply it without assessing its safety to the crop (phytotoxicity).

Because it is not possible to test all material combinations and environmental tank mix combinations, always conduct a test with the intended tank mix on a small area prior to any large scale applications. To determine if a combination is phytotoxic to a specific crop, spray a small area of foliage and fruit, and then evaluate 3-7 days later for visual symptoms.

Always test the compatibility of spray adjuvants (i.e. stickers, spreaders, wetting agents) with Rampart 6.8 before use. Do not use acidifying type compatibility agents.

To determine the compatibility of Rampart 6.8 with other products, use the following jar compatibility test:

1. Add the correct proportions of each product and the appropriate quantity of water to clean container
2. Thoroughly mix
3. Let stand for 3-5 minutes

If the mixture remains in solution or can be remixed readily, the products are considered compatible.

CHEMIGATION

APPLICATIONS THROUGH SPRINKLER AND DRIP IRRIGATION SYSTEMS

Apply this product through microjet, drip (surface and subsurface), solid set, and center pivot sprinkler irrigation systems. Do not apply this product through any other type of irrigation system.

Preparation of Injection Equipment: Remove scale, pesticide residues and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

Application Instructions: Fill tank with $\frac{1}{2}$ to $\frac{3}{4}$ of desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of Rampart 6.8, then the remaining volume of water. Set sprinkler to deliver 0.1 to 0.3 inch of water per acre. Start sprinkler and uniformly inject the solution of Rampart 6.8 into the irrigation water line so as to deliver the desired rate per acre. Inject the Rampart 6.8 solution with a positive displacement pump into the main line ahead of a right angle turn to ensure adequate mixing. For other questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

Note: For Microjet and Drip Irrigation Systems: Avoid further irrigation of the treated area for 24 to 48 hours after treatment has been completed. For Solid Set and Center Pivot Irrigation Systems: Avoid further irrigation of the Rampart 6.8 treated area until after the foliage has completely dried. This will help to prevent washing the chemical off the crop.

GENERAL PRECAUTIONS FOR APPLICATIONS THROUGH SPRINKLER AND DRIP IRRIGATION SYSTEMS

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Maintain continuous agitation in mix tank during mixing and application to assure uniformity solution: Do not apply when wind speed favors drift, when system connections or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled or drained. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from non-uniform distribution of chemigation water.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation shall shut down and adjust the system as needed.

Do not connect an irrigation system (including greenhouse system) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.

FOOD CROPS

Rampart 6.8 fungicide will control Downy Mildew, *Erwinia amylovora*, Pythium, Fusarium, Rhizoctonia and Phytophthora disease (such as purple blotch, late blight, blue mold, soft rot, root rot, brown rot, foot rot, crown/collar rot, canker, fruit/foliar rot, fire blight, heart rot, pink rot, pythium leak, red stele, leather rot) on agronomic crops. Refer to the specific crop groupings below for additional diseases treated and instructions.

BERRY CROPS: Blackberry (including bingberry, black satin berry, boysenberry, Cherokee blackberry, Chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Grapes (all types), Himalayaberry, hullberry, Lavacaberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, nectarberry, ollalieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, youngberry, and varieties and/or hybrids of these), Blueberry, Cane Berries, Cranberry, Currant, Elderberry, Gooseberry, Huckleberry, Loganberry, Rasperry (black and red) and Strawberry.

Use Precautions:

- Do not exceed the highest rate per application.
- Do not apply at less than 3 day intervals.
- Do not apply to plants that are dormant or heat or moisture stressed.
- To avoid undesirable copper phytotoxicity, do not apply Rampart 6.8 foliarly to plants treated with copper based compounds at less than 20 day intervals unless instructed to do so by your consultant.

Application Instructions: Make applications prior to disease development in conjunction with good cultural management practices. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Use rates and frequency are designed to prevent plant injury; use the higher rate when disease pressure is severe. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations.

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium* Rhizoctonia*	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre.	Disease Prevention Program †: Apply lower rate at 2-4 week intervals after plants become established.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	Disease Control Program : Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity.
	Root Dip	2 quarts/100 gallons of water, 2/3 fl. oz./gallon of water.	Apply as pre-plant dip to the roots for 15-30 minutes. Plant within 24 hours after dipping.
	Irrigation	Overhead : 2 to 4 quarts in at least 100 gals. of water. Drip : 2 to 3 qts. in at least 100 gals. of water.	Apply with normal irrigation schedule. Follow Disease Prevention Program or Disease Control Program as stated above.
Downy Mildew	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre.	Apply at 2-3 week intervals.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	Follow Disease Prevention Program or Disease Control Program as stated above.

† If you are unsure these conditions exist consult your local Extension Agent.

*Not approved for this use in CA.

BRASSICA CROPS: Broccoli (all types), Brussels sprouts, Cabbage (all types), Cauliflower, Cavalo broccoli, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach and Rape greens.

Use Precautions:

- Do not exceed the highest rate per application.
- Do not apply at less than 3 day intervals.
- Do not apply to plants that are dormant or heat or moisture stressed.
- Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).
- To avoid undesirable copper phytotoxicity, do not apply Rampart 6.8 foliarly to plants treated with copper based compounds at less than 20-day intervals unless instructed to do so by your consultant.

Application Instructions: Make applications prior to disease development in conjunction with good cultural management practices. Allow foliage to completely dry after application.

Use rates and frequency are designed to prevent plant injury; use the higher rate when disease pressure is severe. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations.

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium* Rhizoctonia*	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre.	Disease Prevention Program †: Apply lower rate at 2-4 week intervals after plants become established.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	Disease Control Program : Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity.
	Root Dip	1 qt./100 gal. of water, 1/3 fl. oz./gal of water	Apply as pre-plant dip immediately prior to planting. (100 gallons is intended to treat the number of slips required to plant one acre.)
	Irrigation	Overhead : 2 to 4 qts. in at least 100 gal. of water. Drip : 2 to 3 qts. in at least 100 gal. of water.	Apply with normal irrigation schedule. Follow Disease Prevention Program or Disease Control Program as stated above.
Downy Mildew	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre.	Apply at 2-3 week intervals.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	Follow Disease Prevention Program or Disease Control Program as stated above.

† If you are unsure these conditions exist consult your local Extension Agent.

*Not approved for this use in CA.

CEREAL GRAINS, FORAGE AND FIBER CROPS: Alfalfa, Barley, Buckwheat, Canola, Clover (all types), Corn (all types), Cotton, Hay, Kudzu, Lespedeza, Lupin, Millet, Oats, Popcorn, Rice, Rye, Safflower, Sainfoin, Sorghum, Sunflower, Teosinte, Trefoil, Triticale, Vetch, Wheat and Wild rice.

Use Precautions:

- Do not exceed the highest rate per application.
- Do not apply at less than 3 day intervals.
- Do not apply to plants that are dormant or heat or moisture stressed.
- Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).
- To avoid undesirable copper phytotoxicity, do not apply Rampart 6.8 foliarly to plants treated with copper based compounds at less than 20-day intervals unless instructed to do so by your consultant.

Application Instructions: Make applications prior to disease development in conjunction with good cultural management practices. Allow foliage to completely dry after application.

Use rates and frequency are designed to prevent plant injury; use the higher rate when disease pressure is severe. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendation.

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium* Rhizoctonia*	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre.	Disease Prevention Program †: Apply lower rate at 2-4 week intervals after plants become established. Disease Control Program: Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.**	Apply with normal irrigation schedule. Follow Disease Prevention Program or Disease Control Program as stated above.
Downy Mildew	Irrigation	Overhead: 2 to 4 qts. in at least 100 gals. of water. Drip: 2 to 3 qts. in at least 100 gals. of water.	Apply at 2-3 week intervals. Follow Disease Prevention Program or Disease Control Program as stated above.
	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre.	
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	

† If you are unsure these conditions exist consult your local Extension Agent.

*Not approved for this use in CA.

**For aerial application in Idaho, Nevada, Oregon and Utah, application may be made in 5 gallons of water/acre.

CITRUS CROPS (Bearing and Non-Bearing) INCLUDING AVOCADO: Avocado, Calamondin, Citrus hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin, Orange (all types), Pummelo, and Satsuma mandarin.

Use Precautions:

- Do not exceed the highest rate per application.
- Do not apply at less than 3 day intervals.
- Do not apply to plants that are dormant or heat or moisture stressed.
- Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).
- To avoid undesirable copper phytotoxicity, do not apply Rampart 6.8 foliarly to plants treated with copper based compounds at less than 20-day intervals unless instructed to do so by your consultant.

Application Instructions: Make applications prior to disease development in conjunction with good cultural practices. Allow foliage to completely dry after application.

Use rates and frequency are designed to prevent plant injury; use the higher rate when disease pressure is severe. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations.

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium* Rhizoctonia*	Foliar	1 to 3 quarts in a minimum of 100 gallons of water/acre.	Disease Prevention Program †: Apply lower rate at 2-4 week intervals after plants become established.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	Disease Control Program: Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity.
	Root Dip	2 qts./100 gal. of water, 2/3 fl. oz./gal. of water.	Apply as pre-plant dip to the roots for 15-30 minutes. Plant within 24 hours after dipping.
	Irrigation	Overhead: 2 to 4 qts. in at least 100 gal. of water. Drip: 2 to 3 qts. in at least 100 gal. of water.	Apply with normal irrigation schedule. Follow Disease Prevention Program or Disease Control Program as stated above.
Downy Mildew	Trunk Injection	3 teaspoons/linear yard of canopy width or 2 in. of trunk diameter at breast height (DBH). Inject directly into tree.	Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications. Repeat 2-4 times a year until control is reached.
	Foliar	1 to 3 quarts in a minimum of 100 gallons of water/acre.	Apply at 2-3 week intervals. Follow Disease Prevention Program or Disease Control Program as stated above.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	

† If you are unsure these conditions exist consult your local Extension Agent.

*Not approved for this use in CA.

CUCURBIT CROPS: Chayote, Chinese waxgourd, Citron melon, Cucumber (all types), Gherkin, Gourd, Muskmelon, Pumpkin, Squash (summer and winter) and Watermelon.

Use Precautions:

- Do not exceed the highest rate per application.
- Do not apply at less than 3 day intervals.
- Do not apply to plants that are dormant or heat or moisture stressed.
- Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).
- To avoid undesirable copper phytotoxicity, do not apply Rampart 6.8 foliarly to plants treated with copper based compounds at less than 20-day intervals unless instructed to do so by your consultant.

Application Instructions: Make applications prior to disease development in conjunction with good cultural management practices. Allow foliage to completely dry after application.

Use rates and frequency are designed to prevent plant injury; use the higher rate when disease pressure is severe. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations.

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium* Rhizoctonia*	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre.	Disease Prevention Program †: Apply lower rate at 2-4 week intervals after plants become established.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	Disease Control Program: Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity.
	Irrigation	Overhead: 2 to 4 qts. in at least 100 gal. of water. Drip: 2 to 3 qts. in at least 100 gal. of water.	Apply with normal irrigation schedule. Follow Disease Prevention Program or Disease Control Program as stated above.
Downy Mildew	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre.	Apply at 2-3 week intervals.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	Follow Disease Prevention Program or Disease Control Program as stated above.

† If you are unsure these conditions exist consult your local Extension Agent.

*Not approved for this use in CA.

HERB AND SPICE CROPS: Allspice, Angelica, Anise, Annatto, Balm, Basil, Borage, Burnet, Camomile, Caper buds, Caraway, Cardamom, Cassia bark and buds, Catnip, Celery seed, Chervil, Chive, Cinnamon, Clary, Clove, Cocoa, Coriander, Costmary, Cilantro, Cumin, Curry, Dill, Fennel, Fenugreek, Grain of paradise, Hops, Horehound, Hyssop, Juniper berry, Lavender, Lemongrass, Lovage, Mace, Marigold, Marjoram, Mint, Mustard, Nasturtium, Nutmeg, Parsley, Pennyroyal, Pepper, Poppy seed, Rosemary, Rue, Saffron, Sage, Savory, Sweet bay, Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff, and Wormwood.

Use Precautions:

- Do not exceed the highest rate per application.
- Do not apply at less than 3 day intervals.
- Do not apply to plants that are dormant or heat or moisture stressed.
- Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).
- To avoid undesirable copper phytotoxicity, do not apply Rampart 6.8 foliarly to plants treated with copper based compounds at less than 20-day intervals unless instructed to do so by your consultant.

Application Instructions: Make applications prior to disease development in conjunction with good cultural management practices. Allow foliage to completely dry after application.

Use rates and frequency are designed to prevent plant injury; use the higher rate when disease pressure is severe. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations.

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium* Rhizoctonia*	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre.	Disease Prevention Program †: Apply lower rate at 2-4 week intervals after plants become established.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	Disease Control Program : Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity.
	Irrigation	Overhead : 2 to 4 qts. in at least 100 gal. of water. Drip : 2 to 3 qts. in at least 100 gal. of water.	Apply with normal irrigation schedule. Follow Disease Prevention Program or Disease Control Program as stated above.
Downy Mildew	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre.	Apply at 2-3 week intervals.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	Follow Disease Prevention Program or Disease Control Program as stated above.

† If you are unsure these conditions exist consult your local Extension Agent.

*Not approved for this use in CA.

LEGUME CROPS: Bean (all types), broad bean (fava bean), Chick pea (garbanzo bean), Guar, Jackbean, Lablab bean, Lentil, Pea (all types), Peanut, Pigeon pea, Soybean and Sword bean.

Use Precautions:

- Do not exceed the highest rate per application.
- Do not apply at less than 3 day intervals.
- Do not apply to plants that are dormant or heat or moisture stressed.
- Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).
- To avoid undesirable copper phytotoxicity, do not apply Rampart 6.8 foliarly to plants treated with copper based compounds at less than 20-day intervals unless instructed to do so by your consultant.

Application Instructions: Make applications prior to disease development in conjunction with good cultural management practices. Allow foliage to completely dry after application.

Use rates and frequency are designed to prevent plant injury; use the higher rate when disease pressure is severe. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations.

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium* Rhizoctonia*	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre.	Disease Prevention Program †: Apply lower rate at 2-4 week intervals after plants become established.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.**	Disease Control Program : Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity.
	Irrigation	Overhead : 2 to 4 qts. in at least 100 gal. of water. Drip : 2 to 3 qts. in at least 100 gal. of water.	Apply with normal irrigation schedule. Follow Disease Prevention Program or Disease Control Program as stated above.
Downy Mildew	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre.	Apply at 2-3 week intervals.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	Follow Disease Prevention Program or Disease Control Program as stated above.

† If you are unsure these conditions exist consult your local Extension Agent.

*Not approved for this use in CA.

**For aerial application in Idaho, Nevada, Oregon and Utah, application may be made in 5 gallons of water/acre.

POME FRUIT CROPS (Bearing and Non-Bearing) INCLUDING PAPAYA: Apple, Crabapple, Loquat, Mayhaw, Papaya, Pear, and Quince.

Use Precautions:

- Do not exceed the highest rate per application.
- Do not apply at less than 3 day intervals.
- Do not apply to plants that are dormant or heat or moisture stressed.
- Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).
- To avoid undesirable copper phytotoxicity, do not apply Rampart 6.8 foliarly to plants treated with copper based compounds at less than 20-day intervals unless instructed to do so by your consultant.

Application Instructions: Make applications prior to disease development in conjunction with good cultural management practices. Allow foliage to completely dry after application.

Use rates and frequency are designed to prevent plant injury; use the higher rate when disease pressure is severe. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations.

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium* Rhizoctonia* Fire Blight (<i>Erwinia amylovora</i>)	Foliar	1 to 3 quarts in a minimum of 100 gallons of water/acre.	Disease Prevention Program [†] : Apply lower rate at 2-4 week intervals after plants become established.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	Disease Control Program : Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity.
	Root Dip	2 qts./100 gal. of water, 2/3 fl. oz./gal. of water	Apply as pre-plant dip to the roots for 15-30 minutes. Plant within 24 hours after dipping.
	Irrigation	Overhead: 2 to 4 qts. in at least 100 gal. of water. Drip: 2 to 3 qts. in at least 100 gal. of water.	Apply with normal irrigation schedule. Follow Disease Prevention Program or Disease Control Program as stated above.
	Trunk Injection	3 teaspoons/linear yard of canopy width or 2 in. of trunk diameter at breast height (DBH). Inject directly into tree.	Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications. Repeat 2-4 times a year until control is reached.
Downy Mildew	Foliar	1 to 3 quarts in a minimum of 100 gallons of water/acre.	Apply at 2-3 week intervals.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	Follow Disease Prevention Program or Disease Control Program as stated above.

[†] If you are unsure these conditions exist consult your local Extension Agent.

*Not approved for this use in CA.

STONE FRUIT CROPS INCLUDING OLIVES: Apricot, Cherry, Nectarine, Olive, Peach, Plum, Plumcot, and Prune.

Use Precautions:

- Do not exceed the highest rate per application.
- Do not apply at less than 3 day intervals.
- Do not apply to plants that are dormant or heat or moisture stressed.
- Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).
- To avoid undesirable copper phytotoxicity, do not apply Rampart 6.8 foliarly to plants treated with copper based compounds at less than 20-day intervals unless instructed to do so by your consultant.

Application Instructions: Make applications prior to disease development in conjunction with good cultural management practices. Allow foliage to completely dry after application.

Use rates and frequency are designed to prevent plant injury; use the higher rate when disease pressure is severe. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations.

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium* Rhizoctonia*	Foliar	1 to 3 quarts in a minimum of 100 gallons of water/acre.	Disease Prevention Program †: Apply lower rate at 2-4 week intervals after plants become established.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	Disease Control Program : Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity.
	Root Dip	2 qts./100 gal. of water, 2/3 fl. oz./gal. of water	Apply as pre-plant dip to the roots for 15-30 minutes. Plant within 24 hours after dipping.
	Irrigation	Overhead: 2 to 4 qts. in at least 100 gal. of water. Drip: 2 to 3 qts. in at least 100 gal. of water.	Apply with normal irrigation schedule. Follow Disease Prevention Program or Disease Control Program as stated above.
	Trunk Injection	3 teaspoons/linear yard of canopy width or 2 in. of trunk diameter at breast height (DBH). Inject directly into tree.	Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications. Repeat 2-4 times a year until control is reached.
Downy Mildew	Foliar	1 to 3 quarts in a minimum of 100 gallons of water/acre.	Apply at 2-3 week intervals.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	Follow Disease Prevention Program or Disease Control Program as stated above.

† If you are unsure these conditions exist consult your local Extension Agent.

*Not approved for this use in CA.

TREE NUT CROPS INCLUDING VINES: Almond, Beech nut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Grapes (all types), Hickory nut, Kiwi, Macadamia nut, Pecan, Pistachio and Walnut.

Use Precautions:

- Do not exceed the highest rate per application.
- Do not apply at less than 3 day intervals.
- Do not apply to plants that are dormant or heat or moisture stressed.
- Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).
- To avoid undesirable copper phytotoxicity, do not apply Rampart 6.8 foliarly to plants treated with copper based compounds at less than 20-day intervals unless instructed to do so by your consultant.

Application Instructions: Make applications prior to disease development in conjunction with good cultural management practices. Allow foliage to completely dry after application.

Use rates and frequency are designed to prevent plant injury; use the higher rate when disease pressure is severe. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations.

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium* Rhizoctonia*	Foliar	1 to 3 quarts in a minimum of 100 gallons of water/acre.	Disease Prevention Program †: Apply lower rate at 2-4 week intervals after plants become established.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	Disease Control Program : Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity.
	Root Dip	2 qts./100 gal. of water, 2/3 fl. oz./gal. of water	Apply as pre-plant dip to the roots for 15-30 minutes. Plant within 24 hours after dipping.
	Irrigation	Overhead: 2 to 4 qts. in at least 100 gal. of water. Drip: 2 to 3 qts. In at least 100 gal. of water.	Apply with normal irrigation schedule. Follow Disease Prevention Program or Disease Control Program as stated above.
	Trunk Injection	3 teaspoons/linear yard of canopy width or 2 in. of trunk diameter at breast height (DBH). Inject directly into tree.	Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications. Repeat 2-4 times a year until control is reached.
Downy Mildew	Foliar	1 to 3 quarts in a minimum of 100 gallons of water/acre.	Apply at 2-3 week intervals.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	Follow Disease Prevention Program or Disease Control Program as stated above.

† If you are unsure these conditions exist consult your local Extension Agent.

*Not approved for this use in CA.

VEGETABLES INCLUDING MINOR FOOD CROPS: (Leafy, Root and Tuber) Amaranth (Chinese spinach, tampala, etc.), Arracacha, Arugula, Arrowroot, Artichoke (all types), Asparagus, Beet (all types), Burdock, Canna, Cardoon, Carrot, Cassava, Celeriac, Celery, Chayote, Chervil (turnip-rooted), Chicory, Chilies, Chrysanthemum (edible), Chufa, Cinchoca, Coconut, Coffee, Corn (all types), Cress (all types), Dandelion, Dasheen, Dock, Eggplant, Endive, Fennel, Garlic, Ginger, Ginseng, Horseradish, Leren, Lettuce (all types), Okra, Onion, Orach, Parsley, Parsnip, Peppers (all types), Pineapple, Purslane, Potato (all types), Radicchio, Radish (all types), Rhubarb, Rutabaga, Salsify (all types), Skirret, Spinach (all types), Sugarcane, Sweet Potato, Swiss Chard, Tanier, Tobacco, Tomato, Turmeric, Turnip, Yam (all types).

Use Precautions:

- Do not exceed the highest rate per application.
- Do not apply at less than 3 day intervals.
- Do not apply to plants that are dormant or heat or moisture stressed.
- Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).
- To avoid undesirable copper phytotoxicity, do not apply Rampart 6.8 foliarly to plants treated with copper based compounds at less than 20-day intervals unless instructed to do so by your consultant.

Application Instructions: Make applications prior to disease development in conjunction with good cultural management practices. Allow foliage to completely dry after application.

Use rates and frequency are designed to prevent plant injury; use the higher rate when disease pressure is severe. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations.

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium* Rhizoctonia*	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre.	Disease Prevention Program †: Apply lower rate at 2-4 week intervals after plants become established.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.**	Disease Control Program : Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity.
	Root Dip	2 qts./100 gal. of water, 2/3 fl. oz./gal. of water	Apply as pre-plant dip immediately prior to planting. (100 gallons is intended to treat the number of slips required to plant one acre.
	Irrigation	Overhead: 2 to 4 qts. in at least 100 gal. of water. Drip: 2 to 3 qts. In at least 100 gal. of water.	Apply with normal irrigation schedule. Follow Disease Prevention Program or Disease Control Program as stated above.
Downy Mildew	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre.	Apply at 2-3 week intervals.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	Follow Disease Prevention Program or Disease Control Program as stated above.
Pink Rot (<i>Phytophthora erythroseptica</i>)	Potato vines: Foliar or Aerial Application	3 to 5 quarts in a minimum of 5 gallons of water/acre.	Apply at 2-3 week intervals. Follow Disease Prevention Program or Disease Control Program as stated above.
Late Blight (<i>Phytophthora infestans</i>) Silver scurf* (<i>Heliniosporium solani</i>) (potatoes only)	Potatoes: In-Storage Application	Humidification, Process and Rinse Waters: Add 0.1 gallon Rampart 6.8 per 0.5 gallons of water.	Humidification, Process and Rinse Waters: Add Rampart 6.8 to water used in humidification, for process water used in post-harvest storage, and for rinses of associated tanks, flumes and lines.

† If you are unsure these conditions exist consult your local Extension Agent.

*Not approved for this use in CA.

**For aerial application in Idaho, Nevada, Oregon and Utah, application may be made in 5 gallons of water/acre.

POST-HARVEST AGRICULTURAL APPLICATION

For control and suppression of labeled post-harvest diseases and storage rots (including Late Blight; Anthracnose; Botrytis; Pink, Brown, Sour, Bitter, Crown, Speckle, Freckle, Finger, Fusarium Rot and other soft rots; Green, Blue and Grey Mold; Black, Chocolate and Cavity Spot) in Potatoes†, Sweet Potatoes†, Yams†, Citrus and citrus hybrids, Avocado†, Carrots†, Pome fruit†, Stone fruit†, Banana† and Plantains†, Mango†, Papaya†, Pineapple†, Lychee†, Mangosteen†, Wax Apple†, Longan†, Carambola†, Guava†, and certain cut flowers†.

†Not registered for use in CA unless accompanied by supplemental labeling

Apply Rampart 6.8 in aqueous applications such as pressure and bin washers or dump and soak tanks. Rampart 6.8 is compatible with most post-harvest chemicals, and will be particularly effective in preventing development of tolerant or resistant strains of fungi when tank mixed with a registered fungicide of a different class of chemistry or when registered chemicals require alternating with a different chemical class. Use a suitable non-ionic surfactant in all post-harvest applications.

†Potatoes, Sweet Potatoes and Yams

DISEASE	APPLICATION RATE	APPLICATION PROGRAM
Suppression** of Late Blight (<i>Phytophthora infestans</i>) and Pink Rot (<i>Phytophthora erythroseptica</i>)	Tuber low volume spray: Apply 0.8 to 1.2 pints Rampart 6.8 per ton of tubers in 0.5 to 1 gallon of water as a spray Dipping: Make up a solution of 1 part Rampart 6.8 to 125 parts water	Tubers should receive uniform, overall coverage of Rampart 6.8 solution. For dips, make sure that tubers are immersed for at least 30 seconds.

†Not registered for use in CA unless accompanied by supplemental labeling

Citrus including, but not limited to Grapefruit, Lemons, Limes, Oranges (sour and sweet), Tangelos, Tangerines, Citrus citron, Citrus hybrids (Mandarin lime, Calamondin, Clementines, Kumquat, Pummelo, and Satsuma mandarin)

DISEASE	APPLICATION RATE	APPLICATION PROGRAM
Suppression** of Brown rot (<i>Phytophthora</i> spp), Green mold (<i>Penicillium digitatum</i>), Blue Mold (<i>Penicillium italicum</i>) and Sour Rot (<i>Geotrichum candidum</i>)	Make up to 1.6% v/v solution of Rampart 6.8 in water (1.6 gallons Rampart 6.8 in 98.4 gallons water). Use approximately 100 gallons of diluted solution per 30 bins* of fruit. For applications to aqueous sprays prior to washing, apply 2.5 pints per 25 gallons spray mixture	Fruit should receive uniform, overall coverage of Rampart 6.8 solution. Make sure fruits are drenched for at least 60 seconds.

*Note: Average bin weight = 800 lbs, or 30 bins weigh 12 Tons.

†For use on Avocado, Pome Fruit (Apples, Pears, Loquat, Mayhaw, Quince), Stone Fruit (Apricots, Cherry (sour and sweet), Nectarine, Peach, Plum, Plumcot, Prune), Banana, Plantain, Mango, Papaya, Pineapple, Guava, Lychee, Longan, Mangosteen Carambola, Wax Apple and Carrots in a Drench Tank or as a Spray

CROP	DISEASE	APPLICATION RATE	APPLICATION PROGRAM
Avocado	Suppression** of Anthracnose (<i>Colletotrichum</i> spp) <i>Dithirella</i> and <i>Cercospora</i> spp.	Apply 3.2 to 4.8 quarts Rampart 6.8 per 100 gallons water. Use approximately 100 gallons of diluted solution per 30 bins* of fruit	Avocado should receive uniform, overall coverage of Rampart 6.8 solution. Replace solution every 30 bins of fruit.
Pome Fruit	Suppression** of Brown Rot (<i>Phytophthora</i> spp.) Blue Mold (<i>Penicillium</i> spp), Bitter Rot (<i>Glomerularia</i> spp) and <i>Monolinia</i> spp.	Apply 3.2 to 4.8 quarts Rampart 6.8 per 100 gallons of water. Use approximately 100 gallons of diluted solution per 30* bins of fruit.	Fruit should receive uniform, overall coverage of Rampart 6.8 solution. Replace solution every 30 bins of fruit.
Stone Fruit	Suppression** of Brown Rot (<i>Phytophthora</i> spp), <i>Monolinia</i> spp, Grey Mold (<i>Botrytis</i>) and Blue mold (<i>Penicillium</i> spp)	Apply 3.2 quarts Rampart 6.8 per 100 gallons of water. Use approximately 100 gallons of diluted solution per 30 bins* of fruit.	Fruit should receive uniform, overall coverage of Rampart 6.8 solution. Replace solution every 30 bins of fruit.
Bananas and Plantains	Suppression** of Crown rot (<i>Fusarium</i> spp.), Anthracnose (<i>Colletotrichum</i> spp), Speckle, Freckle and Finger Rot	Dehanding Tank: Apply 1.6 to 2.4 gallons Rampart 6.8 per 200 gallons of water. Replenish every 2 days Fungicide application. Apply as a spray mix with other fungicides at 0.8 to 1.6 pints Rampart 6.8 per 25 gallons of water	Apply Rampart 6.8 solution to dehanded fruit once it has passed through delatexing tank. Apply Rampart 6.8 as part of a post-harvest fungicide treatment to bananas and plantains
Mangos, Papayas, Pineapple, Lychee, Longan, Carambola, Wax Apple, Mangosteen, Guava	Suppression** of <i>Anthracnose</i> spp, Black Spot, chocolate Spot, <i>cercospora</i> , <i>Alternaria</i> , <i>Botryodiplodia</i> , <i>Phomopsis</i> , <i>Cladosporium</i>	Fruit Dip: make up a 1.6% v/v solution of Rampart 6.8 in water (2 oz Rampart 6.8 plus 126 oz. water to make 1 gallon). Use one gallon of the diluted mix per 4000 lbs of fruit	Fruit should receive uniform, overall coverage of Rampart 6.8 solution. Apply with other fungicides in hot or cold water.
Carrots	Suppression** of Cavity spot, Fusarium Rot, Phytophthora rot, grey Mold (<i>Botrytis cineria</i>), Crown Rot and Black Rot	Apply 2.4 pints Rampart 6.8 per 75 gallons of water. Use one gallon of the diluted mix per 6000 lbs of carrots	Immerse carrots for 5-10 seconds prior to storage. Replace solution when it appears dirty

†Not registered for use in CA unless accompanied by supplemental labeling

* Note: Average Bin Weight = 800 lbs; or 30 bins weigh 12 tons

†For use on Cut Flowers, Roses and Chrysanthemums

CROP	DISEASE	APPLICATION RATE	APPLICATION PROGRAM
Roses, Chrysanthemums, Carnations and Other Cut Flowers	For the Suppression** of post-harvest storage rots, including Botrytis, Anthracnose, vascular wilts and soft rots	Apply 0.4 – 1.2 pints Rampart 6.8 per 100 gallons in the hydrating solution Use the lower rate (0.4 pints / 100 gallons) when low pest pressure is present. Use the higher rate (1.2 pints / 100 gallons) when high pest pressure is present	Cut flower stems should receive uniform, overall coverage of Rampart 6.8 solution when immersing. Do not immerse the flower heads

†Not registered for use in CA unless accompanied by supplemental labeling

** Suppression: For disease suppression, use Rampart 6.8 in combination with another registered bactericide or fungicide registered for the same crops as listed on this label. Note that Rampart 6.8 has suppressive properties on certain diseases caused by bacteria or fungi.

ORNAMENTALS

Rampart 6.8 fungicide will control Pythium, Downy Mildew, Phytophthora, and other labeled diseases on ORNAMENTALS grown in commercial nurseries and greenhouses, golf courses, parks and commercial landscapes.

Rampart 6.8 may be applied to all ornamental and landscape plants including (but not limited to):

Aglaonema	Aphelandra	Arbovitae	Ash
Azaleas	Bougainvilles	Boxwood	Bushes
Cattleya sinneri	Cedars	Ceanothus	Cotoneaster
Cissus	Diffenbachia	Dogwood	Ferns
Ficus	Flowering Trees	Hibiscus	Hard Wood Trees
Holly	Home Lawns	Juniper	Magnolia Trees
Maple	Pines	Oaks	Palms
Peperomia	Photinia	Pittosporum	Philodendron
Pieris	Pothos	Red woods	Rhododendron
Roses	Schefflera	Sedum	Sempervivum
Shrubs	Spathiphyllum	Sycamore	Taxus media
Zygocactus			

Use Precautions:

- Do not exceed the highest rate per application.
- Do not apply at less than 3 day intervals.
- Do not apply to plants that are dormant or heat or moisture stressed.
- To avoid undesirable copper phytotoxicity, do not apply Rampart 6.8 foliarly to plants treated with copper based compounds at less than 20-day intervals unless instructed to do so by your consultant.

Application Instructions:

Apply Rampart 6.8 prior to disease development in conjunction with good cultural management practices. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Use rates and frequency are designed to prevent plant injury; use the higher rate when disease pressure is severe. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations.

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium* Rhizoctonia* Fire Blight (<i>Erwinia amylovora</i>) Bacterial Blight* (<i>Xanthomonas campestris</i>) Green Wild (<i>Ralstonia</i> spp.)	Foliar Spray	1 to 2 qt./100 gal of water or 1/3 fl. oz. to 2/3 fl. oz per gal. of water	Disease Prevention Program †: Apply lower rate at 2-4 week intervals after plants become established. Disease Control Program : Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity.
	Soil Drench	12-24 fl. oz./100 gals. of water or 1/8 to ¼ fl. oz. per gallon of water	Apply 1-2 pts. of diluted solution per sq. foot. Adjust volume as needed to thoroughly wet media. Apply at 2-3 week intervals and repeat as needed.
	Soil Incorporation	1-2 pts./cubic yard of soil mix	Mix 1-2 pints of Rampart 6.8 with enough water to wet one cubic yard of soil; mix immediately before potting. If conditions are favorable for disease development, it is recommended to make additional applications as a foliar spray or soil drench.
	Irrigation	Overhead : 1 to 2 qts./100 gallons of water Drip : 1 to 2 qts./100 gallons of water	Apply with normal irrigation schedule. Follow Disease Prevention Program or Disease Control Program as stated above.
	Trunk Injection	3 teaspoons/linear yard of canopy width or 2 in. of trunk diameter at breast height (DBH). Inject directly into tree.	Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications. Repeat 2-4 times a year until control is reached.
	Root Dip	2 qts./ 100 gal. of water or 2/3 fl. oz./gal. of water	Dip transplants into diluted solution of Rampart 6.8. Thoroughly wet plant root mass immediately before transplanting. Keep roots submerged for 1-2 minutes.
Downy Mildew	Foliar Spray	1-2 qts./ 100 gal. of water or 1/3 fl. oz. to 2/3 fl. oz./gallon of water	Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed. Follow Disease Prevention Program or Disease Control Program as stated above.

† If you are unsure these conditions exist consult your local Extension Agent.

*Not approved for this use in CA.

BEDDING PLANTS

Rampart 6.8 fungicide will control Downy Mildew, Pythium, Phytophthora and other labeled diseases on bedding plants grown in commercial nurseries and greenhouses, golf courses, parks and commercial landscapes.

Rampart 6.8 may be applied to all bedding plants including (but not limited to) the following:

Ageratum	Algerian ivy	Alyssum	Anthurium
Artemisia	Aster	Baby's Breath	Begonia
Caladium	Carnation	Chrysanthemum	Coleus
Columbine	Daisy	Delphinium	Easter Lily
English Ivy	Foxglove	Gaillardia	Geranium
Gloxinia	Impatiens	Marigold	Pansy
Petunia	Phlox	Pinks	Poinsetta
Primrose	Prostrate	Rosemary	Salvia
Snapdragon	Verbena	Vinca	Zinnia

Use Precautions:

- Do not exceed the highest rate per application.
- Do not apply at less than 3 day intervals.
- Do not apply to plants that are dormant or heat or moisture stressed.
- To avoid undesirable copper phytotoxicity, do not apply Rampart 6.8 foliarly to plants treated with copper based compounds at less than 20-day intervals unless instructed to do so by your consultant.

Application Instructions:

Apply Rampart 6.8 at 2-3 week intervals prior to disease development in conjunction with good cultural management practices. Apply diluted solution to thoroughly wet foliage and allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Use rates and frequency are designed to prevent plant injury; use the higher rate when disease pressure is severe. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations.

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium* Rhizoctonia* Fire Blight (<i>Erwinia amylovora</i>)	Foliar Spray	1-2 qt./100 gal of water or 1/3 fl. oz. to 2/3 fl. oz per gal. of water	Disease Prevention Program †: Apply lower rate at 2-4 week intervals after plants become established. Disease Control Program : Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity
	Root Dip	2 qts./100 gals. of water or 2/3 fl. oz./gallon of water.	Dip transplants into diluted solutions of Rampart 6.8. Thoroughly wet plant root mass immediately before transplanting. Keep roots submerged for 1 to 2 minutes.
	Soil Drench	12-24 fl. oz./100 gals. of water or 1/8 to ¼ fl. oz. per gallon of water	Apply 1-2 pts. of diluted solution per sq. foot. Adjust volume as needed to thoroughly wet media. Apply at 2-3 week intervals and repeat as needed.
	Irrigation	Overhead : 1 to 2 qts. in at least 100 gal. of water. Drip : 1 to 2 qts. in at least 100 gal. of water.	Apply with normal irrigation schedule. Follow Disease Prevention Program or Disease Control Program as stated above.
Downy Mildew	Foliar Spray	1 to 2 qts./ 100 gal. of water (1/3 fl. oz. to 2/3 fl. oz./gallon of water)	Follow Disease Prevention Program or Disease Control Program as stated above.

† If you are unsure these conditions exist consult your local Extension Agent.

*Not approved for this use in CA.

CONIFERS AND TREES IN COMMERCIAL NURSERIES, PLANTATIONS AND FORESTS, LANDSCAPES AND PARKS (INCLUDING CHRISTMAS TREES, OAKS AND PINES) APPLICATIONS BY COMMERCIAL APPLICATORS

Rampart 6.8 applied in conjunction with good cultural management practices will control Phytophthora root rot in conifers including (but not limited to) Douglas fir, spruce and pines as well as other Phytophthora disease of Oaks (Sudden Oak Death) and forest trees.

Use Precautions:

- Do not apply to trees that are under heat or moisture stress.
- Do not graze livestock in treated conifer nurseries and/or plantations.
- Do not feed forage to livestock from treated nurseries and/or plantations.

Application Instructions:

Use the higher rates when disease pressure is severe.

Disease	Application Method	Rate	Application Program
Phytophthora	Foliar Spray	1-2 qt./100 gal of water	Disease Prevention Program †: Apply lower rate at 2-4 week intervals after plants become established. Disease Control Program : Apply higher rate at 2-3 week intervals until control is reached.
		or 1/3 fl. oz. to 2/3 fl. oz per gal. of water. Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed.	
	Soil Drench	12 to 24 fl. oz./100 gals. of water or 1/8 to ¼ fl. oz. per gallon of water	Apply 1 to 2 pts. of diluted solution per sq. foot. Adjust volume as needed to thoroughly wet media. Apply at 2 to 3 week intervals and repeat as needed.
	Trunk Injection	3 teaspoons/linear yard of canopy width or 2 in. of trunk diameter at breast height (DBH). Inject directly into tree.	Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications.
	Root Dip	2 qts/ 100 gal. of water or 2/3 fl. oz./gal. of water	Apply as a pre-plant dip to the roots for 15-30 minutes. Plant within 24 hours after dipping.

† If you are unsure these conditions exist consult your local Extension Agent.

TURF

(INCLUDES RESIDENTIAL AND COMMERCIAL LAWN APPLICATIONS BY COMMERCIAL APPLICATORS)

Rampart 6.8 fungicide will control Pythium blight, Summer Complex (Bermuda and others), Dollar Spot, Grey Leaf Spot, Brown Patch, Anthracnose, and damping-off diseases in turf grasses on sod farms and lawns, commercial turf production, golf courses (including tees and greens), parks and commercial landscapes.

Use Precautions:

- Do not graze animals on treated turf.
- Do not feed clippings from treated turf to livestock and poultry.

Application Instructions:

Begin preventive applications when conditions first favor disease and repeat at recommended intervals. Use higher rates under severe disease conditions. Do not mow or irrigate treated areas until sprays have completely dried.

Disease	Application Method	Rate	Application Program
Pythium (Yellow Turf)	Foliar Spray	2 to 3 fl. oz./1000 sq. ft.	Apply amount in minimum of 2 gallons of water per 1000 sq. ft. Apply at 10 to 21 day intervals as needed.
Rhizoctonia* Sclerotinia* Helminthosporium* Alternaria*		1 to 3 quarts/100 gallons of water/acre.	For sod farms and large areas: Apply diluted solution to thoroughly wet foliage. Apply at 2 to 3 weeks intervals and repeat as needed.
Anthracnose* (<i>Colletotrichum gram-inicola</i>)	Irrigation	Overhead: 2 to 3 qts./ acre in at least 100 gal. of water	Apply with normal irrigation schedule. Apply at 10 to 21 day intervals and repeat as needed.

*Not approved for this use in CA.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Nonrefillable Container (five gallons or less): Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

Nonrefillable Container (greater than five gallons): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

FORMULATED FOR LOVELAND PRODUCTS, INC.
P.O. BOX 1286, GREELEY, COLORADO 80632-1286

INTENTIONALLY LEFT BLANK