

**PRODUCT INFORMATION SUPPLEMENTAL LABEL  
TO ADD NEW CROP USES IN CALIFORNIA**



EPA Reg. No. 71512-3-279

**Active Ingredient:**

Cyazofamid\* .....34.5%  
**Other Ingredients:** ..... 65.5%  
100.0%

\*4-chloro-2-cyano-N,N-dimethyl-5-(4-methylphenyl)-1H-imidazole-1-sulfonamide (CA)  
 Contains 3.33 pounds Cyazofamid Per Gallon (400 grams per liter)

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED.**

This Supplemental labeling must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for Ranman Fungicide before applying. Carefully follow all precautionary statements and application use directions. Use of Ranman Fungicide according to this supplemental labeling is subject to all use precautions and limitations imposed by the labeling affixed to the container for Ranman Fungicide.

Crop	Diseases	Use Rate Fl. Oz Product Per Acre (lb. ai/A)	Instructions
Carrot	Cavity spot ( <i>Pythium ultimum</i> ) Root Dieback ( <i>P. violae</i> , <i>P. sulcatum</i> ) Forking ( <i>P. irregulare</i> , <i>P. splendens</i> )	6 (0.156)	<p><b>Resistance Management:</b> DO NOT apply more than 5 sprays of Ranman per crop. Alternate sprays of Ranman with a fungicide with a different mode of action.</p> <p><b>Application instructions:</b> Pre-plant incorporated (broadcast or band): Apply in sufficient water to obtain adequate coverage within 3 days of planting and mechanically till into the soil to a depth of at least 2 inches or incorporate with at least 1/4 inch of water. Surface applications (broadcast or band): Subsequent applications may be made beginning at 14 days after plant emergence and continue on a 14-21 day schedule. Apply in sufficient water to obtain adequate coverage with the applications directed to the base of the plant. Ranman should be incorporated into the soil with 1/2 to 1 inch of water. If irrigation is not immediately available after the application, then the application should be made in sufficient water to allow penetration into the soil. Ranman may be applied via any overhead irrigation system. Follow directions outlined in the Application and Calibration Techniques For Sprinkler Irrigation section of the label. Ranman 400SC should be applied during the last 2 hours of the irrigation cycle to allow for adequate soil penetration. For banded applications a 6 to 8 inch band is recommended (See formula to calculate amount required in the band). Calculate the amount of Ranman needed for band treatments by the formula:</p> $\frac{\text{band width in inches}}{\text{row spacing in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$ <p><b>Restrictions:</b> DO NOT use more than 30 fl oz per growing season. DO NOT use any adjuvant when applying to carrots. DO NOT apply within 14 days of harvest. Crops on this label may be planted immediately after the last treatment. Do not plant other crops not registered for this product within 30 days after the last application.</p>



FMC Corporation  
 Agricultural Products Group  
 1735 Market Street  
 Philadelphia PA 19103