



Belchim Crop Protection US Corporation
 2751 Centerville Road, Suite 100
 Wilmington, DE 19808
 855-445-7990
www.belchimusa.com

FIFRA 2(ee) Recommendation FOR THE USE OF RELOAD® FUNGICIDE FOR THE SUPPRESSION OF DAMPING OFF AND ROOT ROTS IN LEAFY VEGETABLE CROPS

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin must be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF ARIZONA.

RELOAD® FUNGICIDE
 EPA Reg. No. 87865-1

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

CROP	DISEASE	RATE pint/acre	USE RECOMMENDATIONS
LEAFY VEGETABLES (Except Brassica Leafy Vegetables) All members of Crop Group 4 including: Amaranth, Arugula (Roquette), Cardoon, Celery, Chinese Celery, Celtuce, Chervil, Chrysanthemum (Edible leaved and garland), Corn salad, Cress (Garden and Upland), Dandelion, Dock (Sorrel), Endive (Escarole), Florence Fennel, Lettuce (Head and Leaf), Orach, Parsley, Purslane (Garden and Winter), Radicchio (Red chicory), Rhubarb, Spinach, New Zealand Spinach, Vine Spinach, Swiss chard	Damping Off and Root Rots <i>(Phytophthora and Pythium spp.)</i>	2.0 – 5.0	Apply in a minimum of 10 gallons of water per acre to obtain good coverage. Begin application when conditions favor disease development and continue on a 7- to 21-day spray interval. Use higher rate and shorter spray interval during times of moderate to high disease pressure. NOTE: DO NOT exceed 7 applications per season.

For Chemical Emergency,
 Spill, Leak, Fire, Exposure or Accident,
 Call CHEMTREC Day or Night
 Within USA and Canada 1-800-424-9300

Reload® is a registered trademark of Belchim Crop Protection USA, LLC