

# Product Bulletin



Loveland Products, Inc., P.O. Box 1286, Greeley, Colorado 80634

## Reaper® 0.15 EC

EPA Reg. No. 34704-923

2(ee) Recommendation<sup>†</sup>

Control of Additional Pest in Florida

### ATTENTION

- <sup>†</sup>This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA or state lead pesticide agency.
- Read and follow all applicable directions for use, precautions and limitations on the federally registered label attached to the product container.

**Additional pest on avocados controlled within existing rate range of the federally registered product label for Reaper:**

Avocados	
Additional Target Pest	Instructions
Mirids ( <i>Daghbertus fasciatus</i> , <i>Daghbertus olivaceous</i> , <i>Rhinacloa</i> sp.)	Apply 10-20 fl. oz. Reaper 0.15 EC per acre. For best results, apply when immature mirids are first observed. Apply in the January to April time period. See restrictions below.

### Restrictions:

- The user must refer to the federally approved **Reaper** label and read and follow all applicable directions for use, restrictions, and precautions.
- Do not apply within 14 days of harvest.
- Ground and aerial application permitted
- Wait at least 30 days before repeating application
- Do not make more than 2 applications of Reaper 0.15 EC or any other foliarly-applied abamectin-containing product per season.
- Do not apply more than 40 fl. oz. per acre (or 0.047 pounds active ingredient per acre) of Reaper 0.15 EC or any other foliarly applied abamectin-containing product per season.
- Do not apply more than 20 fl. oz. per acre (or 0.023 pounds active ingredient per acre) of Reaper 0.15 EC or any other foliarly applied abamectin-containing product per application.

- Do not allow livestock to graze in treated orchards.

The user must refer to the federally approved Reaper® 0.15 EC label and read and follow all applicable directions for use, restrictions, and precautions.

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This recommendation should be in the possession of the user at the time of pesticide application.