

# SUPPLEMENTAL LABEL

(Contains additional information, which may not appear on package label.)

**RESTRICTED USE PESTICIDE**  
**TOXIC TO FISH, MAMMALS, AND AQUATIC ORGANISMS**  
**FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR**  
**THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.**

## Reaper 0.15 EC

EPA REG. NO 34704-923

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE PACKAGE LABEL ARE TO BE FOLLOWED.**

**THIS LABEL MUST BE IN POSSESSION OF USER AT THE TIME OF APPLICATION.**

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**Cotton:** Reaper 0.15 EC is an emulsifiable concentrate containing 0.15 lb abamectin/gal. Which, when mixed with water according to the Directions for Use, will control spider mites (twospotted, carmine, Pacific and strawberry) on cotton.

For control of spider mites on cotton, mix with water and apply at the rates given in the following table:

Pest	Fl. oz./A		Instructions
	Mid-season/Lay-by	Early Season	
Spider Mites Twospotted Carmine Pacific Strawberry	8 – 16	4 – 6	Mix with water and begin application when mites are first noticed and repeat if necessary. Do not use less than 4 fl. oz./A <b>Early Season Application:</b> Apply only when cotton is less than 10 inches tall. Use 4 – 6 fl. oz./A; do not use less than 4 fl. oz./A. <b>Mid-Season/lay-by application:</b> Use 8 – 16 fl. oz./A depending on size of the plant and density of the foliage.

#### Use Restrictions – Cotton

**Location:** Do not apply to cotton east of the Mississippi River

**PHI:** Do not apply within 20 days of harvest

#### Application Method:

**Early Season:** Use only ground application equipment; aerial application is not permitted.

**Mid-Season/lay-by:** Application with ground application equipment and aircraft (fixed-winged or helicopter) is permitted in a minimum of 5 gals of water/A.

**Amount of Water:** Do not apply in fewer than 5 gallons of water/A with aerial application equipment.

**Application Interval:** Wait at least 21 days before repeating application.

**Maximum Amount per Season:** Do not apply more than 32 fl. oz./A or more than 2 applications per growing season.

**Grazing Restriction:** Do not feed cotton foliage to livestock or allow livestock to graze treated cotton.

Packaged for:



P.O. Box 1286  
Greeley, CO 80632