



TECHNICAL INFORMATION BULLETIN
FOR THE USE OF REGALIA® Rx ON
CEREAL GRAINS

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(EE) RECOMMENDATIONS ARE RECOGNIZED.

REGALIA®

EPA Reg. No. 84059-3

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CROPS	DISEASES	APPLICATION METHOD	RATES	USE RECOMMENDATIONS
Cereal Grains Barley Buckwheat Rye Grain Amaranth* Oat Millets Rice *	<u>Powdery Mildew</u> <u>(Erysiphe graminis)</u> <u>Leaf Spots</u> <u>(Dreschlera,</u> <u>Cochliobolus,</u> <u>Cercospora)</u> <u>Rust (Puccinia spp.)</u> <u>Septoria</u> <u>Leaf/Speckled</u> <u>Leaf Spot/Blotch</u> <u>(Septoria spp.)</u> <u>Smut (Tilletia spp.)</u> <u>Tan Spot</u> <u>(Pyrenophora</u> <u>tritici- repentis)</u>	<u>Foliar</u> <u>(Ground)</u> <u>Disease</u> <u>Applications</u>	1-2 quarts per acre	For ground applications for disease control, apply this product in 15–40 gallons of water per acre. It is important to apply this product at the flag leaf stage to maximize yield. Apply this product preventatively or when the first disease symptoms appear. Repeat applications in 7-14 day intervals depending upon crop growth and disease pressure. When plants are under high disease pressure, tank mix this product with another fungicide for more effective control.
		<u>Foliar</u> <u>(Aerial)</u> <u>Disease</u> <u>Applications</u>	1 quart per acre	For aerial applications, apply this product in a minimum of 5 gallons of water per acre. It is important to apply this product at the flag leaf stage to maximize yield. Apply this product preventatively or when the first disease symptoms appear. Repeat applications in 7-14 day intervals depending upon crop growth and disease pressure. When plants are under high disease pressure, tank mix this product with another fungicide for more effective control.

Cereal Grains Barley Buckwheat Rye Grain Amaranth* Oat Millets Rice *		<u>Foliar (Ground)</u> <u>Plant Health Applications</u>	1 pint per acre	For ground applications to maximize yields, apply this product in 15-40 gallons of water per acre. Apply with a preferred tank mix partner at 50% - 100% Flag Leaf emergence to maximize yields.
		<u>Foliar (Aerial)</u> <u>Plant Health Applications</u>	1 pint per acre	For aerial applications to maximize yields, apply this product in a minimum of 5 gallons of water per acre. Apply with a preferred tank mix partner at 50% - 100% Flag Leaf emergence to maximize yields.
	<ul style="list-style-type: none"> • <i>Fusarium</i> spp. • <i>Phytophthora</i> spp. • <i>Pythium</i> spp. • <i>Rhizoctonia</i> spp. • <i>Verticillium</i> spp. 	SEED TREATMENT applications	1.5–2.5 fluid ounces per 100 lbs. seed	For suppression of soil-borne diseases, apply this product as a seed treatment at the rate of 1.5–2.5 fl. oz. per 100 lbs. seed.
* Not for use in California				

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.

This 2(ee) recommendation expires December 31, 2019.

Regalia[®] and Regalia[®] Rx are registered trademarks of Marrone Bio Innovations, Inc.
 © Copyright Marrone Bio Innovations, Inc., 2009

Marrone Bio Innovations, Inc.
 1540 Drew Ave.
 Davis, California 95618
 Telephone (530) 750-2800
 www.marronebioinnovations.com