



TECHNICAL INFORMATION BULLETIN
FOR THE USE OF REGALIA® Rx ON
COTTON

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(EE) RECOMMENDATIONS ARE RECOGNIZED.

REGALIA®

EPA Reg. No. 84059-3

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CROPS	DISEASES	APPLICATION METHOD	RATES	USE RECOMMENDATIONS
COTTON	Alternaria Leaf Spot, Boll Rot (<i>Alternaria</i> spp.) Anthracnose, Boll Rot (<i>Glomeria</i> spp.) Ascochyta Blight, Boll Rot (<i>Ascochyta</i> spp.) Cercospora Blight and Leaf Spot (<i>Cercospora</i> spp.)	Foliar (Ground) Disease Applications	1-2 quarts per acre	For ground applications for foliar and boll rot disease control, apply this product preventatively in 15–40 gallons of water per acre prior to disease development using sufficient volume for thorough coverage. Repeat applications at 7–14 day intervals as necessary to maintain control.
	Diplodia Boll Rot (<i>Diplodia</i> spp.) Hard Lock, Boll Rot (<i>Fusarium</i> spp.) Leaf Spot (<i>Corynespora cassicola</i>) Phoma Blight, Boll Rot (<i>Phoma</i> spp.) Rust (<i>Puccinia</i> spp.), (<i>Phykopsora</i> spp.) Stemphylium Leaf Spot (<i>Stemphylium</i> spp.)	Foliar (Aerial) Disease Applications	0.5 - 1 quart per acre	For aerial applications, apply this product in a minimum of 3 gallons of water per acre.

COTTON	<i>Fusarium</i> spp. <i>Phytophthora</i> spp. <i>Pythium</i> spp. <i>Rhizoctonia</i> spp. <i>Verticillium</i> spp.	In-Furrow applications	1–4 quarts per acre or 2.2–8.8 fluid ounces per 1000 ft. row	For in-furrow applications, at planting apply this product as an in-furrow spray at the rate of 1–4 quarts per acre or 2.2–8.8 fluid ounces per 1000 feet of row according to the chart in the SOIL TREATMENT USE DIRECTIONS section. Apply this product in 5–15 gallons of water so as the spray is directed into the seed furrow just before the seeds are covered.
		Seed Treatment applications	7.5–15 fluid ounces per 100 lbs. seed	For suppression of soil borne diseases, apply this product as a seed treatment at the rate of 7.5–15 fl. oz per 100 lbs. seed.
COTTON*		Foliar (Ground) Plant Health Applications*	1 pint per acre	For ground applications to maximize yields, apply this product in 15-40 gallons of water per acre.
		Foliar (Aerial) Plant Health Applications*	1 pint per acre	For aerial applications to maximize yields, apply this product in a minimum of 5 gallons of water per acre.
* Not for Use in California				

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.

This 2(ee) recommendation expires December 31, 2019.

Regalia[®] and Regalia[®] Rx are registered trademarks of Marrone Bio Innovations, Inc.
 © Copyright Marrone Bio Innovations, Inc., 2009

Marrone Bio Innovations, Inc.
 1540 Drew Ave.
 Davis, California 95618
 Telephone (530) 750-2800
www.marronebioinnovations.com