



TECHNICAL INFORMATION BULLETIN  
FOR THE USE OF REGALIA® Rx ON  
PEANUT

**FIFRA Section 2(ee) Recommendation**

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(EE) RECOMMENDATIONS ARE RECOGNIZED.

**REGALIA®**

**EPA Reg. No. 84059-3**

**DIRECTIONS FOR USE**

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

| CROPS  | DISEASES   | APPLICATION METHOD                               | RATES               | USE RECOMMENDATIONS  |
|--------|--|--|---------------------|--|
| PEANUT | Aspergillus Crown Rot ( <i>Aspergillus niger</i> )   | <a href="#">Foliar Disease Applications</a>      | 1-4 quarts per acre | For foliar applications, apply this product preventatively in 20–50 gallons of water per acre.<br><br>Repeat applications at 7–14 day intervals.<br><br>Tank-mix this product with another fungicide labeled for the target disease. |
|        | Damping-off ( <i>Aspergillus flavus</i> ), ( <i>Fusarium</i> spp.), ( <i>Pythium</i> spp.), ( <i>Rhizoctonia</i> spp.)<br><br>Rhizoctonia Foliar Blight, Peg, and Root Rot ( <i>Rhizoctonia solani</i> )<br><br>White Mold ( <i>Sclerotium rolfsii</i> ) | <a href="#">Foliar Plant Health Applications</a> | 1 pint per acre     | For ground applications to maximize yields, apply this product in 15-40 gallons of water per acre.   |

|               |   |                             |  |   |
|---------------|---|-----------------------------|--|---|
| <b>PEANUT</b> | <i>Aspergillus</i> Crown Rot<br><i>(Aspergillus niger)</i><br><i>Fusarium</i> spp.<br><i>Phytophthora</i> spp.<br><i>Pythium</i> spp.<br><i>Rhizoctonia</i> spp.<br><i>Verticillium</i> spp.<br>White Mold<br><i>(Sclerotium rolfsii)</i> | In-Furrow applications      | 1–4 quarts per acre or 2.2–8.8 fluid ounces per 1000 ft. row | For in-furrow applications at planting, apply this product as an in-furrow spray at the rate of 1–4 quarts per acre or 2.2–8.8 fluid ounces per 1000 feet of row according to the chart in the SOIL TREATMENT USE DIRECTIONS section. Apply this product in 5–15 gallons of water so as the spray is directed into the seed furrow just before the seeds are covered.   |
|               |   | Soil Drench Applications    | 1–3 quarts per 100 gallons of water                          | For soil drench applications, apply this product at a concentration of 1–3 quarts per 100 gallons of water, and at a sufficient rate to thoroughly soak the growing media and root zone. Make an initial application of this product during or shortly after transplant to reduce transplant shock, suppress soil-borne diseases and improve root growth. Multiple drench applications can be made on a 10-14-day interval. |
|               |   | Seed Treatment Applications | 1.5–2.5 fluid ounces per 100 lbs. seed                       | For suppression of soil-borne diseases, apply this product as a seed treatment at the rate of 1.5–2.5 fl. oz per 100 lbs. seed.   |

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.**

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

**This recommendation must be in the possession of the user at the time of pesticide application.**

This 2(ee) recommendation expires December 31, 2019.

Regalia<sup>®</sup> and Regalia<sup>®</sup> Rx are registered trademarks of Marrone Bio Innovations, Inc.  
 © Copyright Marrone Bio Innovations, Inc., 2009

Marrone Bio Innovations, Inc.  
 1540 Drew Ave.  
 Davis, California 95618  
 Telephone (530) 750-2800  
[www.marronebioinnovations.com](http://www.marronebioinnovations.com)