



TECHNICAL INFORMATION BULLETIN
FOR THE USE OF REGALIA® Rx ON
NON-GRASS ANIMAL FEED

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(EE) RECOMMENDATIONS ARE RECOGNIZED.

REGALIA®

EPA Reg. No. 84059-3

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CROPS	DISEASES	APPLICATION METHOD	RATES	USE RECOMMENDATIONS
Non-grass Animal Feed Alfalfa (except California) Clover Kudzu Lespedeza Lupin Sainfoin Trefoil Vetch	Alfalfa Wilt (<i>Xylella</i> spp.) Anthracnose and Black Stem Rot (<i>Colletotrichum trifolii</i>) Bacterial Wilt (<i>Clavibacter michiganense</i>) Downy Mildew (<i>Peronospora trifoliorum</i>) Fusarium wilt (<i>Fusarium oxysporum</i>) Leafspots and Blotches (<i>Pseudopeziza medicaginus</i> , <i>Stemphyllium</i> spp., <i>Cercospora</i> spp., <i>Stagonospora</i> spp.) Mosaic viruses Powdery Mildew (<i>Erysiphe polygoni</i>) Sclerotinia stem and crown rot (<i>Sclerotinia sclerotiorum</i>)	Foliar (Ground) <u>Disease Applications</u>	1-2 quarts per acre	For ground applications to optimize disease control and to maximize yields, apply this product in 15–40 gallons of water per acre. It is important to apply this product at the flag leaf stage to maximize yield. Apply this product preventatively or when the first disease symptoms appear. Repeat applications on 7–14 day intervals depending upon crop growth and disease pressure. When the plants are under high disease pressure, tank-mix this product with another fungicide for more effective control.
	Seedling and Damping Off Disease Complex, including Root and Crown Rots (<i>Pythium</i> , <i>Phytophthora</i> , <i>Rhizoctonia</i> , and <i>Stagonospora</i> spp.) Spring Black Stem (<i>Phoma medicaginus</i>) Verticillium wilt (<i>Verticillium albo-atrum</i>)	Foliar (Aerial) <u>Disease Applications</u>	0.5 - 1 quart per acre	For aerial applications, apply this product in a minimum of 5 gallons of water per acre. It is important to apply this product at the flag leaf stage to maximize yield. Apply this product preventatively or when the first disease symptoms appear. Repeat applications on 7-14 day intervals depending upon crop growth and disease pressure. When plants are under high disease pressure, tank mix this product with another fungicide for more effective control.



Non-grass Animal Feed Alfalfa (except California) Clover Kudzu Lespedeza Lupin Sainfoin Trefoil Vetch		Foliar (Ground) Plant Health Applications	1 pint per acre	For ground applications to maximize yields, apply this product in 15-40 gallons of water per acre. For plant health applications, apply 10-14 days prior to cutting.
		Foliar (Aerial) Plant Health Applications	1 pint per acre	For aerial applications to maximize yields, apply this product in a minimum of 5 gallons of water per acre. For plant health applications, apply 10-14 days prior to cutting.

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.

This 2(ee) recommendation expires December 31, 2019.

Regalia® and Regalia® Rx are registered trademarks of Marrone Bio Innovations, Inc.
 © Copyright Marrone Bio Innovations, Inc., 2009

Marrone Bio Innovations, Inc.
 1540 Drew Ave.
 Davis, California 95618
 Telephone (530) 750-2800
www.marronebioinnovations.com