



TECHNICAL INFORMATION BULLETIN
FOR THE USE OF REGALIA® Rx IN AERIAL APPLICATIONS

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROPS EXIST AND WHERE 2(Ee) RECOMMENDATIONS ARE RECOGNIZED.

REGALIA® Rx
EPA Reg. No. 84059-3

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CROP	APPLICATION METHOD	RATE*	USE RECOMMENDATIONS
Cereal Grains (barley, sorghum, rice, wheat) Corn (field corn, popcorn, seed corn, silage corn, sweet corn) Cotton Oil Seed Crops (canola, sunflower) Peanut Sugar Beets Sugarcane Soybean Tobacco	Foliar (Aerial)	1 pint per acre*	Applying Regalia® Rx has been shown to increase leaf chlorophyll content and increase soluble protein content in some crops. These effects often lead to improved crop quality and/or yields. For aerial applications, apply this product in 2–5 gallons of water per acre. Mixing Regalia® Rx with other labeled fungicides has been shown to enhance control of targeted diseases. If compatibility with another product is unknown, a jar test should be conducted.

*Plant health uses only. For disease control rates, follow instructions on EPA-registered label.

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.

REGALIA® is a registered trademark of Marrone Bio Innovations, Inc.
© Copyright Marrone Bio Innovations, Inc., 2014