

Bayer Environmental Science A division of Bayer CropScience LP 5000 CentreGreen Way, Suite 400 Cary, NC 27513 1-866-99BAYER (1-866-992-2937)

Rejuvra™

For control of Stinknet (globe chamomile) in rangelands, CRP lands and natural areas.

EPA Reg. No. 432-1609

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF ARIZONA, NEVADA AND NEW MEXICO

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

Timing of Application

Rejuvra may be applied at any time of the year. For pre-emergence control of annual weeds apply several weeks prior to weed seed germination. For maximum weed control, apply during periods when sufficient precipitation to activate the herbicide is expected prior to target weed germination, but avoid application if heavy rain is expected which can move treated soil into areas with crops or other desirable vegetation. For applications targeting emerged or established weeds, mix Rejuvra with appropriate post-emergence herbicides (see Herbicide Mixtures section).

Use Rates

The desired rate of Rejuvra depends on the residual weed activity required and restrictions on the maximum amounts listed in the table below.

Ilee Rates

The desired rate of Rejuvra depends on the residual weed activity required and restrictions on the maximum amounts listed in the table below.

Rejuvra Rates for Rangeland, CRP and Natural Areas

	Rate Range (fl oz/A)	Maximum Single Use Rate (fl oz/A)	Maximum Allowed in a 12-Month Period (fl oz/A)
Areas grazed by livestock and/or rotated for hay	3.5** - 5	5	6
Areas not grazed by live- stock and not cut for hay	3.5** - 7	7	10

^{**} The 3.5 fl oz rate of Rejuvra should only be used when weed pressure is low and when less preemergence residual control is desired. DO NOT harvest hay within 40 days of any single application.

IMPORTANT

Refer to the Rejuvra™ and any tank-mix partner labels for information on adjuvants; weed species controlled; additional use directions; rotations crop intervals and precautions. The most restrictive provision on any label should apply.

For <u>MEDICAL</u> And <u>TRANSPORTATION</u> Emergencies <u>ONLY</u> Call 24 Hours A Day 1-800-334-7577. For <u>PRODUCT USE</u> Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions. For additional information call toll-free 1-866-99BAYER (1-866-992-2937).