


US81761388C (140430) LABMC REQUIEM EC 2.5 gallon US extended text label  
colors: cmyk 7/18/14



# REQUIEM<sup>®</sup>

EC



**INSECTICIDE AND ACARICIDE FOR CONTROL OR SUPPRESSION OF THRIPS, WHITEFLIES, PLANT-FEEDING MITES, LEAFMINERS, AND LEAF HOPPERS IN AGRICULTURAL CROPS.**

**NOT FOR RESIDENTIAL USE**

**ACTIVE INGREDIENT:**  
Terpene Constituents of the Extract of  
*Chenopodium ambrosioides* near *ambrosioides*  
as Synthetically Manufactured ..... 16.75%  
**OTHER INGREDIENTS:** ..... 83.25%  
Total: 100.00%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

FOR ADDITIONAL PRECAUTIONARY STATEMENTS:  
See Inside Booklet.

**EPA Reg. No. 264-1159      EPA Est. No. 264-MEX-001**

USE OF PRODUCT INDICATES ACCEPTANCE OF  
"CONDITIONS FOR SALE AND WARRANTY"

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY**  
Call 24 Hours A Day 1-800-334-7577  
For **PRODUCT USE** Information Call 1-800-331-2867

Produced for:  
Bayer CropScience LP  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
REQUIEM is a registered trademark of Bayer.  
©2014 Bayer CropScience  
Product of Mexico

140430C 07/14  
US81761388C

5.75"

#### FIRST AID

<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.</b>	

#### PRECAUTIONARY STATEMENTS

##### HAZARD TO HUMANS AND DOMESTIC ANIMALS

##### CAUTION

Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear the appropriate Personal Protective Equipment (PPE).

##### PERSONAL PROTECTIVE EQUIPMENT

Applicators, mixers, loaders and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Shoes plus socks
- Protective eyewear
- Coveralls for high-pressure handwand and groundboom applicators.
- For overhead exposure wear chemical resistant headgear.
- For greenhouse uses, applicators and other handlers must wear a NIOSH-approved respirator with an organic vapor (OV) cartridge or canister with any R, P or HE filter.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls: When handlers use enclosed cabs in a manner that meets requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

5.5625"

5.75"

#### USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. For terrestrial uses: Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not apply when weather conditions favor drift or runoff from treated areas. Do not apply when a rain event is expected within 2 hours. Groundboom and air-blast applications must maintain a spray buffer of 15 feet from any water body. Aerial applications must maintain a spray buffer of 400 feet from any water body. See the Directions for Use for details.

#### DIRECTIONS FOR USE

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

READ THE ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH THE LABEL. FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR INSECT CONTROL, AND/OR ILLEGAL RESIDUES. For use only as described on this label. Not for isolation or deformulation.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the State or Tribal agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural insecticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Shoes plus socks

**SHAKE WELL BEFORE USE**

5.5625"

5.75"

**GENERAL INFORMATION**

REQUIEM® EC is a contact insecticide/acaricide for use in the control or suppression of many foliage feeding pests including thrips, plant-feeding mites, whiteflies, leafminers, and leaf hoppers infesting labeled crops. The emulsifiable concentrate of REQUIEM EC must be mixed with water and applied as a foliar spray with equipment equipped for conventional insecticide spraying.

REQUIEM EC is toxic to aquatic invertebrates. Users must maintain the buffer zones specified in the application instruction sections for ground and aerial applications. Do not apply when a rain event is expected within 2 hours.

Do not apply this product through any type of irrigation system.

IMPORTANT: READ CONDITIONS FOR SALE AND WARRANTY BEFORE USE

**APPLICATION INSTRUCTIONS**

**General**

Use REQUIEM EC early in the pest cycle when local thresholds are reached or before insects reach damaging levels. REQUIEM EC is a contact insecticide and thorough coverage is necessary for optimum control. Use appropriate nozzles and nozzle configurations, along with sufficient spray volume to achieve thorough coverage. Rates of REQUIEM EC may need to increase with increased insect populations and/or large plant canopies. It is not necessary to use a surfactant with REQUIEM EC.

When using REQUIEM EC with a new crop, test on a small area before applying to the entire field. Do not apply REQUIEM EC more than 10 times per crop production cycle.

**Spray Drift**

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower/treatment coordinator are responsible for considering all of these factors when making decisions. Where states have more stringent regulations, they should be observed. Refer to the Aerial Drift Reduction Information section of this label for additional directions and precautions applicable to both ground and aerial applications.

**Ground**

This product can be applied by commonly used ground equipment, such as groundboom, hose-end, pressurized, air-blast, greenhouse and hand-held sprayers. Consult spray nozzle and accessory catalogues for specific information on proper equipment calibration. Maintain agitation during mixing and application to assure uniform product suspension. Thorough coverage of all foliage is essential for effective pest control or suppression. Use the application rate indicated for the appropriate crop in the Application Rate tables of this label in sufficient water to achieve thorough coverage. Overall, to achieve good coverage, use proper spray pressure, gallonage per acre, nozzles, nozzle spacing and ground speed.

Groundboom and air-blast spray applicators must not apply within 15 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds). Groundboom and air-blast applicators must also account for wind direction and speed. Only apply this product if the wind direction favors on-target deposition. Do not apply when the wind velocity exceeds 10 mph.

Use of air-blast spray equipment may result in overhead exposure and requires the use of chemical-resistant headgear.

5.5625"

5.75"

**Aerial**

This product can be applied by aerial application. Use the application rate indicated for the appropriate crop in the Application Rate tables of this label in sufficient water to achieve thorough coverage, typically between 2 – 20 gallons of water per acre depending upon the crop. Aerial applicators must not apply within 400 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

**MIXING INSTRUCTIONS**

Shake container well before use. REQUIEM EC must be diluted with water for spray applications. Partially fill the spray tank with clean water and begin agitation. Add the specified amount of REQUIEM EC to the tank. Finish filling the tank to the proper volume to obtain the desired application rate per acre as specified in the crop and pest tables listed below. It is critical that the spray solution be agitated during mixing and application to ensure a uniform suspension. Do not allow spray mixture to stand overnight or for prolonged periods. When applying REQUIEM EC via ground application, use dilutions no more concentrated than 2% v/v.

**Compatibility:** Do not combine REQUIEM EC in the spray tank with other pesticides, surfactants, adjuvants, or fertilizers if there has been no previous experience or use of the combination to show it is physically compatible, and non-injurious under your use conditions.

Do not use REQUIEM EC in tank mixes or in a program or in rotation with products containing captan (with REQUIEM EC either preceding or following captan applications) in any crop. Use of REQUIEM EC with captan-containing products may result in phytotoxicity.

Do not use REQUIEM EC in tank mixture combinations or in rotation with any product containing a label warning against using surfactants, methylated seed oils, oils or other adjuvants.

As with many Emulsifiable Concentrate (EC) formulations, certain combinations can exacerbate phytotoxicity problems. Products which are phytotoxic when used alone may be more phytotoxic when used with an EC formulation such as REQUIEM EC. Bayer CropScience has not tested all varieties and cultivars of all crops and potential tank-mix or rotational combinations.

To ensure crop safety, test REQUIEM EC in combination and in rotation with other products on a small portion of the crop. Wait at least 24-72 hours to evaluate results before applying to the entire field. If phytotoxicity is noted, and if it is determined that the level of phytotoxicity will result in economic losses to the crop, do not make the application to the crop.

To ensure physical compatibility of tank-mix combinations they must be evaluated prior to use. When products are physically compatible, a homogeneous solution is formed. To determine the physical compatibility of this product with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to approximately one quart of water with agitation. Add dry formulations first, then flowables, then emulsifiable concentrates last. After thorough mixing, allow this mixture to stand for 15 minutes. If the mixture in the test jar remains uniform for 15 minutes, the combination is compatible and can be used. If separation occurs (e.g. oils float to top, clumps of solids form, etc.), the combination is incompatible, do not use the mixture. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

5.5625"

5.75"

#### **Use of Adjuvants**

The use of adjuvants with REQUIEM EC is not required.

#### **Biological Control Agents**

This product is compatible with the following biological control agents: *Encarsia formosa*, *Amblyseius fallacis*, *Phytoseiulus persimilis*, *Orius insidiosus*, *Aphidius colemani*, *Zetzellia mali*, *Neoseiulus fallacis*, *Typhlodromus pyri*, *Stethorus punctum*, lacewings, syrphid fly larvae, phytoseiid mite predators, aphid midges, lady beetles, and other predacious mites. Wait at least 2 hours after application before introducing new biological agents, or spray the evening before the day of introduction of new biological control agents.

Bayer CropScience is committed to the practice of sound resistance management programs that include rotation and tank mixing with other treatments having a variety of modes of actions.

#### **PREHARVEST INTERVAL**

REQUIEM EC can be applied up to and including the day of harvest. However, note that the restricted entry interval (REI) is 4 hours.

#### **AERIAL DRIFT REDUCTION INFORMATION**

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower/treatment coordinator are responsible for considering all of these factors when making decisions. Where states have more stringent regulations, they should be observed. Note: This section is advisory in nature and does not supersede the mandatory label requirements.

All application equipment must be properly maintained and calibrated using appropriate carriers.

#### **Information on Droplet Size**

When used for aerial applications, the best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets will reduce drift potential but will not prevent drift if applications are made improperly or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

#### **Controlling Droplet Size**

Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure - Do not exceed the nozzle manufacturer's specified pressures. For many nozzle types, lower pressure produces larger droplets. When high flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of Nozzles - Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation - Orienting nozzles, so that the spray is released parallel to the airstream, produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

5.5625"

5.75"

**Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid-stream nozzles that are oriented straight back produce the largest droplets and the lowest drift. Use medium or coarser spray according to the ASAE 572 definition for standard nozzles or VMD for spinning atomizer nozzles.

**Boom Width**

For aerial applications using fixed-wing aircraft, the boom width must not exceed 75% of the wingspan. For aerial applications using helicopters, the boom width must not exceed 90% of the rotary blade.

**Application Height**

Do not release spray at a height greater than 10 feet above the top of the ground or the crop canopy unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind. If application includes a no-spray zone, do not release spray at a height greater than 10 feet above ground or canopy.

**Swath Adjustment**

Use upwind swath displacement. When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the upwind and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

**Wind**

Apply only when wind speed is 3 - 10 miles per hour as measured by an anemometer. Drift potential is lowest between wind speeds of 3 – 10 miles per hour. Many factors, however, including droplet size and equipment type, determine drift potential at any given speed. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

**Temperature and Humidity**

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

**Temperature Inversions**

Do not apply during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light, variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

5.5625"

5.75"

**Sensitive Areas**

This pesticide should only be applied when the potential for drift to adjacent, sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas). Do not allow spray to drift from the application site and contact people, structures people occupy at any time and the associated property, parks and recreation areas, nontarget crops, aquatic and wetland areas, woodlands, pastures, rangelands, or animals. Applicator must maintain the buffer zones specified in the application instruction sections for ground and aerial applications.

**FOR USE ON THE FOLLOWING CROPS**

**(REQUIEM EC has a 0-Day PreHarvest Interval (PHI) for all crops contained on this label. However, note that the restricted entry interval (REI) is 4 hours.)**

**BRASSICA (COLE) LEAFY VEGETABLES**

Including Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage (Bok Choy and Napa), Cauliflower, Cavalo, Collards, Kale, Kohlrabi, Mustard Greens, Rape Greens, and all other Brassica leafy vegetable crops.

**Pests and Application Rates:**

Pests	REQUIEM EC (qts/acre)	Remarks
Whiteflies: • Silverleaf whitefly	2 - 4	Apply before pests reach damaging levels. Repeat applications as needed to maintain control. Do not wait until heavy populations have become established. Use the high rate when conditions are favorable for heavy pressure.

**BULB VEGETABLES**

Including Garlic, Leek, Onion (dry bulb, green, and Welch), and Shallot and other bulb vegetable crops.

**Pests and Application Rates:**

Pests	REQUIEM EC (qts/acre)	Remarks
Thrips: • Onion thrips • Western flower thrips • Tobacco thrips	1.5 - 3	Begin applications as soon as thrips are seen in the crop. Continue applications as needed. Thrips will seek shelter in hard-to-reach parts of the plant. Use insecticide nozzles with sufficient water volume and spray pressure for thorough coverage.

5.5625"



5.75"

**CITRUS FRUITS**

Including Grapefruit, Lemon, Lime, Orange, Tangerine, Citrus hybrids, Pummelo, and other citrus fruit crops.

**DO NOT APPLY TO CITRUS FRUITS BY AERIAL APPLICATION**

**Pests and Application Rates:**

Pests	REQUIEM EC (qts/acre)	Remarks
Asian citrus psyllid* Mites: <ul style="list-style-type: none"> <li>• Citrus red mite</li> <li>• Citrus rust mite</li> <li>• Six-spotted spider mite</li> <li>• Texas citrus mite</li> <li>• Two-spotted spider mite</li> </ul>	2 - 4	Treat when pests appear or in accordance with local economic thresholds. Repeat applications as needed to maintain control. The rate of REQUIEM EC will depend on tree size and pest pressure. Use a lower rate for light infestation and/or small trees and a higher rate for heavy infestations and/or large trees. When applying to large trees, use higher water volumes to ensure thorough coverage.

**\*NOT REGISTERED FOR USE IN CALIFORNIA ON ASIAN CITRUS PSYLLID**

**CUCURBIT VEGETABLES**

Including Cucumber, Citron melon, Edible Gourds, Gherkin, Muskmelons (including hybrids and/or cultivars of Cucumis melo), Pumpkin, Summer and Winter Squash, Watermelon, Cantaloupe, and other cucurbit vegetable crops.

**Use of REQUIEM EC in tank mixes with some products including, but not limited to, those containing strobilurins or chlorothalonil may result in phytotoxicity. Do not apply products containing chlorothalonil within 4 days after application of REQUIEM EC. Do not apply REQUIEM EC for 30 days following an application of chlorothalonil.**

**Pests and Application Rates:**

Pests	REQUIEM EC (qts/acre)	Remarks
Thrips: <ul style="list-style-type: none"> <li>• Melon thrips</li> <li>• Western flower thrips</li> </ul>	2 - 3	Begin applications as soon as thrips are seen in the crop. Continue applications as needed. Thrips will seek shelter in hard-to-reach parts of the plant. Use appropriate nozzles with sufficient water volume and spray pressure for thorough coverage.
Whiteflies: <ul style="list-style-type: none"> <li>• Greenhouse whitefly</li> <li>• Sweetpotato whitefly</li> <li>• Silverleaf whitefly</li> </ul>		Apply before pests reach damaging levels. Repeat applications as needed to maintain control. Do not wait until heavy populations have become established. Use the high rate when conditions are favorable for heavy pressure.

5.5625"

5.75"

**FRUITING VEGETABLES**

Including Eggplant, Ground Cherry, Pepino, Pepper, Tomatillo, Tomato, and other fruiting vegetable crops.

**Pests and Application Rates:**

Pests	REQUIEM EC (qts/acre)	Remarks
Thrips: <ul style="list-style-type: none"> <li>• Western flower thrips</li> <li>• Chili thrips</li> <li>• Melon thrips</li> <li>• Florida flower thrips</li> <li>• Eastern flower thrips</li> </ul>	2 - 4	Begin applications as soon as thrips are seen in the crop. Continue applications as needed. Thrips will seek shelter in hard-to-reach parts of the plant. Use appropriate nozzles with sufficient water volume and spray pressure for thorough coverage.
Whiteflies: <ul style="list-style-type: none"> <li>• Silverleaf whitefly</li> </ul>	2 - 3	Apply before pests reach damaging levels. Repeat applications as needed to maintain control. Do not wait until heavy populations have become established. Use the high rate when conditions are favorable for heavy pressure.
Leafminers ( <i>Liriomyza</i> spp.)	3 - 4	For Suppression: Begin treatments before thresholds have been reached.

5.5625"

**GRAPE**

**Do not use REQUIEM EC on grapes east of the Rockies. Use of REQUIEM EC on grapes after fruit set can diminish waxy bloom.**

**DO NOT APPLY TO GRAPES BY AERIAL APPLICATION**

**Pests and Application Rates:**

Pests	REQUIEM EC (qts/acre)	Remarks
Leafhoppers: <ul style="list-style-type: none"> <li>• Grape leafhopper</li> <li>• Variegated leafhopper</li> </ul>	2 - 4	For Suppression: Begin treatments before thresholds have been reached.
Mites: <ul style="list-style-type: none"> <li>• Pacific spider mites</li> <li>• Willamette spider mite</li> <li>• Two-spotted spider mite</li> </ul>		Begin applications when thresholds have been reached. The rate of REQUIEM EC will depend on plant size and pest pressure. Use a lower rate for light infestation and/or small plant canopy and a higher rate for heavy infestations and/or large plants. Adjust equipment, spray volume, and spray pressure to ensure thorough coverage of infested parts of the crop.

5.75"

**LEAFY VEGETABLES**

Including Lettuce, Arugula, Celery, Spinach, Parsley, Radicchio, Florence Fennel, Romaine, Swiss Chard, and other leafy vegetable crops.

**Pests and Application Rates:**

Pests	REQUIEM EC (qts/acre)	Remarks
Whiteflies: • Silverleaf whitefly	2 - 3	Apply before pests reach damaging levels. Repeat applications as needed to maintain control. Do not wait until heavy populations have become established. Use the high rate when conditions are favorable for heavy pressure.
Leafminers ( <i>Liriomyza</i> spp.)	3 - 4	For Suppression: Begin treatments before thresholds have been reached.

**OKRA\***

**\*NOT REGISTERED FOR USE ON OKRA IN CALIFORNIA**

**Pests and Application Rates:**

Pests	REQUIEM EC (qts/acre)	Remarks
Thrips: • Melon thrips • Western Flower thrips • Chili thrips • Florida Flower thrips	2 - 4	Begin applications as soon as thrips are seen in the crop. Continue applications as needed. Thrips will seek shelter in hard-to-reach parts of the plant. Use insecticide nozzles with sufficient water volume and spray pressure for thorough coverage.
Whiteflies		Apply before pests reach damaging levels. Repeat applications as needed to maintain control. Do not wait until heavy populations have become established. Use the high rate when conditions are favorable for heavy pressure.

**ROOT / TUBER AND CORM VEGETABLES**

Including Carrot, Potato, Sweet Potato, Cassava, Beets, Ginger, Horseradish, Radish, Ginseng, Turnip, True Yam, and other root/tuber and corm crops.

**Pests and Application Rates:**

Pests	REQUIEM EC (qts/acre)	Remarks
Psylla: • Potato psyllid*	2 - 3	Begin treatments when thresholds have been reached.

**\*NOT REGISTERED FOR USE ON POTATO PSYLLID IN CALIFORNIA**

5.5625"

5.75"

### STONE FRUITS

Including Apricot, Cherry, Nectarine, Peach, Plum, Plumcot, Prune, and other stone fruit crops

**Do not use of REQUIEM in tank mixes with chlorothalonil. Do not apply products containing chlorothalonil or captan within 30 days after application of REQUIEM. Do not apply REQUIEM for 30 days following an application of chlorothalonil or captan.**

#### DO NOT APPLY TO STONE FRUITS BY AERIAL APPLICATION

##### Pests and Application Rates:

Pests	REQUIEM EC (qts/acre)	Remarks
Mites: <ul style="list-style-type: none"><li>• Two-spotted spider mite</li></ul>	2 - 4	Begin applications when thresholds have been reached. The rate of REQUIEM EC will depend on plant size and pest pressure. Use a lower rate for light infestation and/or small plant canopy and a higher rate for heavy infestations and/or large plant canopy. Adjust equipment, spray pressure, and spray volumes to ensure thorough coverage.

### TREE NUTS

Including Almond, Beech nut, Cashew, Chestnut, Filbert, Hickory nut, Pecan, Pistachio, Walnut, and other tree nut crops.

**Do not use of REQUIEM EC in tank mixes with chlorothalonil. Do not apply products containing chlorothalonil within 30 days after application of REQUIEM EC. Do not apply REQUIEM EC for 30 days following an application of chlorothalonil.**

#### DO NOT APPLY TO TREE NUTS BY AERIAL APPLICATION

##### Pests and Application Rates:

Pests	REQUIEM EC (qts/acre)	Remarks
Mites: <ul style="list-style-type: none"><li>• Pacific spider mite</li><li>• European red mite</li><li>• Two-spotted spider mite</li><li>• Strawberry spider mite</li></ul>	2 - 4	Begin applications when thresholds have been reached. The rate of REQUIEM EC will depend on plant size and pest pressure. Use a lower rate for light infestation and/or small plant canopy and a higher rate for heavy infestations and/or large plant canopy.

5.5625"

5.75"

### OILSEED CROPS

Canola, Castor, Cotton, Flax, Rapeseed, Safflower, Sesame, Sunflower, and other oilseed crops (including those grown for seed or oil production).

#### Pests and Application Rates:

Pests	REQUIEM EC (qts/acre)	Remarks
Mites: • Two-spotted spider mite	0.5 – 2	Begin applications when thresholds have been reached. The rate of REQUIEM EC will depend on plant size and pest pressure. Use a lower rate for light infestation and/or small plant canopy and a higher rate for heavy infestations and/or large plant canopy. Adjust equipment, spray pressure, and spray volumes to ensure thorough coverage.
Whiteflies: • Silverleaf whitefly	2 – 4	

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

#### Pesticide Storage

Store in a cool dry place. Avoid freezing.

#### Pesticide Disposal

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry). Do not contaminate water when disposing of equipment washwater or rinsate. Pesticide wastes may be toxic. Improper disposal of unused pesticide, washwater or rinse water is a violation of federal law.

#### Container Disposal

Non-refillable container. Do not reuse or refill this container.

Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

5.5625"

5.75"

**IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product at once for a refund of the purchase price.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties, and Limitations of Liability. These terms may only be modified by a written document signed by a duly authorized representative of Bayer CropScience LP.


**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LP. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

Bayer

5.5625"



5.75"

Lot No.:

Net Contents: 2.5 Gal (9.46 L)

.6875"

# REQUIEM<sup>®</sup>

EC

**INSECTICIDE AND ACARICIDE FOR CONTROL OR SUPPRESSION OF THRIPS, WHITEFLIES, PLANT-FEEDING MITES, LEAFMINERS, AND LEAF HOPPERS IN AGRICULTURAL CROPS.**

**NOT FOR RESIDENTIAL USE**

**ACTIVE INGREDIENT:**

Terpene Constituents of the Extract of *Chenopodium ambrosioides* near *ambrosioides* as Synthetically Manufactured ..... 16.75%  
OTHER INGREDIENTS: ..... 83.25%  
Total: 100.00%

EPA Reg. No. 264-1159 EPA Est. No. 264-MEX-001

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

FOR ADDITIONAL PRECAUTIONARY STATEMENTS:  
See Attached Booklet.

For **MEDICAL** And **TRANSPORTATION** Emergencies  
**ONLY** Call 24 Hours A Day 1-800-334-7577  
For **PRODUCT USE** Information Call 1-800-331-2867

**FIRST AID**

<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>

**In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.**

**PRECAUTIONARY STATEMENTS**  
**HAZARD TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**

Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear the appropriate Personal Protective Equipment (PPE).

**FOR ADDITIONAL PRECAUTIONARY STATEMENTS:**

See attached label booklet: Personal Protective Equipment, User Safety Recommendations, and Environmental Hazards

**DIRECTIONS FOR USE: See attached booklet.**

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage**

Store in a cool dry place. Avoid freezing.

**Pesticide Disposal**

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry). Do not contaminate water when disposing of equipment washwater or rinsate. Pesticide wastes may be toxic. Improper disposal of unused pesticide, washwater or rinse water is a violation of federal law.

**Container Disposal**

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

Bayer CropScience LP  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
REQUIEM is a registered trademark of Bayer.  
©2014 Bayer CropScience

140430C 07/14  
US81761388C

6.75"

5.5625"

.5"

6.75"