



<b>FIFRA 2(ee) Recommendation</b>
EPA Reg. No. 279-3588
<b>FOR DISTRIBUTION AND USE ONLY IN CALIFORNIA</b>
<b>THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.</b>

This label for Rhyme™ fungicide is valid until December 5, 2024, or until withdrawn, canceled or suspended.

<b>IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.</b>
<b>THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.</b>

### For Management of Botryosphaeria panicle and shoot blight and Phomopsis canker in Walnuts (including black and English (Persian) walnuts)

<b>Directions For Use</b>				
Apply <u>Rhyme</u> fungicide in sufficient water volume to ensure thorough coverage as a foliar application. Apply preventatively or when conditions are favorable for disease development.				
<b>Crop</b>	<b>Disease</b>	<b>Rate (fl oz/A)</b>	<b>Application Directions</b>	<b>Retreatment Interval (Days)</b>
Walnut	Botryosphaeria panicle and shoot blight Phomopsis canker	7	Apply preventatively and repeat as necessary on a 7-day interval when conditions are favorable for disease development.	7

**RESTRICTIONS:**

- No single application may exceed 7 fl oz of product/A.
- Do not apply more than 4 applications per year.
- Do not apply more than 28 fl oz of product/A per year.
- Do not apply more than 0.455 lb ai flutriafol/A per year.
- Restricted Entry Interval (REI) is 12 hours
- Preharvest Interval (PHI) – Do not apply within 14 days of harvest

R-4357 120619 12-09-16