

**THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.**



EPA Reg. No. 279-3588

**2(ee) Recommendation**

**FOR DISTRIBUTION AND USE ONLY IN THE STATES OF CT, GA, NJ, MA, MI, MD, NC, OH, PA, SC, VA, WV**

**THIS FIFRA SECTION 2(EE) EXPIRES JANUARY 30, 2021**

To permit the use of Rhyme<sup>®</sup> fungicide in peach as a foliar application for control of rusty spot (*Podosphaera leucotricha*).

**Directions For Use**

- IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.
- ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.
- This recommendation must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for Rhyme<sup>®</sup> Fungicide before applying.

**Crop**

- Peach

**Rates and Timing**

Diseases	Product Rate (fl. oz./A)	Application Directions	Application Interval (Days)
Rusty spot ( <i>Podosphaera leucotricha</i> )	7 (0.113 lb ai/A)	Start applications at petal fall and continue through 3rd Cover (3C) or until terminal shoot growth ceases and shoots harden off or conditions are no longer favorable for disease development	7

**Restrictions:**

- No single application may exceed 7 fl. oz. of product/A.
- Do not apply more than 4 applications per year.
- Do not apply more than 28 fl. oz. of product/A per year.
- Do not apply more than 0.455 lb. ai. of flutriafol/A per year.
- Restricted Entry Interval (REI) is 12 hours.
- Preharvest Interval (PHI): Do not apply within 7 days of harvest.



FMC Corporation  
2929 Walnut Street  
Philadelphia, PA 19104

FMC and Rhyme are trademarks of FMC Corporation  
Copyright © 2018