
PRODUCT INFORMATION



Syngenta Crop Protection, Inc.
Greensboro, NC 27409
www.syngenta-us.com

**SUPPLEMENTAL LABEL FOR RIDOMIL GOLD® COPPER
FUNGICIDE
EPA Reg. No. 100-804**

To revise Pre Harvest Interval (PHI) for grapes

Active Ingredients:

Mefenoxam (CAS No. 70630-17-0).....	4.8%
Related Compounds.....	0.2%
Copper Hydroxide*: (Cu(OH) ₂)	60.0%
Other Ingredients:	35.0%
Total:	100.0%

*Metallic Copper equivalent.....39.1%

KEEP OUT OF REACH OF CHILDREN.

DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

Before using Ridomil Gold Copper as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling supersedes the Directions for Use on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

GRAPES

Use Ridomil Gold/Copper as a foliar, post-bloom application for control of downy mildew of grapes caused by *Plasmopora viticola*.

Apply 1 pack (5 lbs. product)/2.5 acres of Ridomil Gold/Copper. Make up to 4 applications beginning before bloom; do not make an application within 42 days of harvest. For late season downy mildew control, apply other registered fungicides.

Notes: (1) To avoid possible illegal residues, do not apply within 42 days of harvest. (2) You may make up to 4 applications of Ridomil Gold/Copper per crop. (3) If other formulations of Ridomil Gold are applied, do not apply more than a total of 0.40 lbs. a.i./acre Ridomil Gold per crop per season when using a combination of formulations containing the Ridomil Gold active ingredient. (4) Do not use on copper sensitive varieties (see local state and extension recommendations for varietal selection).

Ridomil Gold® Copper and the Syngenta logo are trademarks of a Syngenta Group Company

© 2004 Syngenta

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP 804B-S3 0704

RID GOL COP 804B-S3 0704 nsp 7-29-04