
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 804B-S4 0211

Ridomil Gold® Copper

Fungicide

For the control of certain diseases in various fruits and vegetables

Active Ingredients:

Mefenoxam*	5.0%
Copper Hydroxide**: (Cu(OH) ₂)	60.0%
<hr/>	
<i>Other Ingredients:</i>	35.0%
Total:	100.0%

*CAS Nos. 70630-17-0 and 69516-34-3

**Metallic Copper equivalent.....39.1%

Ridomil Gold Copper is a wettable powder packaged in a water-soluble bag.

KEEP OUT OF REACH OF CHILDREN.

DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

EPA Reg. No. 100-804

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Ridomil Gold Copper as permitted according to this Supplemental Labeling, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application.

syngenta

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CANEBERRY (BLACKBERRIES AND RASPBERRIES) SUBGROUP*

* Blackberry subgroup includes: Andean blackberry, arctic blackberry, bingleberry, black satin berry, boysenberry, brombeere, California blackberry, Cherokee blackberry, Chesterberry, Cheyenne blackberry, common blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, evergreen blackberry, Himalayaberry, hullberry, lavacaberry, loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, mora, mures deronce, nectarberry, Northern dewberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, Southern dewberry, tayberry, youngberry, and zarzamora.

Ridomil Gold Copper, when used as a foliar spray in a preventive disease control program, provides control of downy mildew of berries.

Apply 1 pack (5 lb. product)/2.5 acres (0.1 lb. a.i./A of mefenoxam) of Ridomil Gold Copper. Begin applications when conditions are favorable for disease, but before infection, and again in 7 days.

Notes: (1) Ridomil Gold Copper may be applied the day of harvest (0-day PHI). (2) Do not make more than 2 applications of a Ridomil Gold prepack (Ridomil Gold Copper or Ridomil Gold Bravo®) per crop. (3) Do not apply with an adjuvant. (4) Do not exceed the equivalent of 1.8 lb. a.i./A per crop of soil-applied and 0.2 lb. a.i./A per crop foliar-applied mefenoxam-containing products, for a maximum seasonal application rate of 2.0 lb, a.i./A.

ONIONS - DRY BULB*, GREEN**, AND ONIONS GROWN FOR SEED

*Includes garlic, onions (dry bulb), and shallots (dry bulb), as well as Great-headed garlic, Serpent garlic, Lily bulb, Chinese onion bulb, Pearl onion, Potato onion bulb.

**Includes green onions, leeks, spring onions or scallions, Japanese bunching onions, green shallots, or green eschalots, as well as chive (fresh leaves), Chinese chive (fresh leaves), Hosta elegans, Fritillaria (leaves), kurrat, lady's leek, Beltsville bunching onions, onion (fresh), macrostem onion, tree tops onion, Welsh onion (tops), shallot (fresh leaves).

Ridomil Gold Copper provides effective control of downy mildew caused by *Peronospora destructor* when used as a foliar spray in a preventive disease control program.

Apply 1 pack of Ridomil Gold Copper (5 lb. product)/2.5 acres in sufficient water to obtain thorough coverage. Begin applications when conditions are favorable for disease, but before infection, and continue at 14-day intervals until the threat of disease is over. Use a suitable spreader-sticker at rates recommended on the product label.

NOTES: (1) Bulb onions: Do not exceed the equivalent of 1.0 lb. a.i./A per crop of soil-applied and 0.4 lb. a.i./A per crop of foliar-applied mefenoxam-containing products. (2) Green onions: Do not exceed the equivalent of 1.0 lb. a.i./A per crop of soil-applied and 0.3 lb. a.i./A per crop of foliar-applied mefenoxam-containing products. (3) Bulb onions: Do not apply within 10 days of harvest (10-day PHI). (4) Green onions: Do not apply within 7 days of harvest (7-day PHI).

SNAP BEANS

Use Ridomil Gold Copper as a foliar application for control of downy mildew caused by *Phytophthora phaseoli*, Pythium pod rot, and Pythium cottony leak. Apply 1 pack (5 lb. product)/2.5 acres (0.1 lb. a.i./A of mefenoxam) of Ridomil Gold Copper. Begin applications at the onset of disease and continue on a 7-day schedule. Do not make more than 2 applications per season.

Notes: (1) Do not apply Ridomil Gold Copper within 7 days prior to harvest (7-day PHI). (2) Do not apply more than 0.4 lb. a.i. mefenoxam/A of Ridomil Gold Copper per season. (3) Do not use an adjuvant. (4) Do not exceed the equivalent of 0.5 lb. a.i./A per crop of soil-applied mefenoxam-containing products at planting and 0.2 lb. a.i./A per crop of foliar-applied mefenoxam-containing products.

SPINACH

Ridomil Gold Copper will control white rust and downy mildew on spinach when applied foliarly following an at-planting application of Ridomil Gold SL at 1-2 pt./acre (0.50-1 lb. a.i./acre).

Apply 1 pack of Ridomil Gold Copper (5 lb. product)/2.5 acres 21 days after the Ridomil Gold SL at-planting application or immediately after each repeated cutting. Make 1 or 2 applications of Ridomil Gold Copper at 14-day intervals, depending on cultural practices. Avoid late-season applications when plants reach full maturity or begin senescence.

Notes: (1) Make up to 2 applications of Ridomil Gold Copper per crop. (2) Do not use an adjuvant. (3) If Ridomil Gold Copper use is planned, apply Ridomil Gold SL either preplant or at planting. **Do not make supplemental applications of Ridomil Gold SL.** (4) If other pesticides containing mefenoxam and/or metalaxyl are also used, the total amount of mefenoxam plus metalaxyl from Ridomil Gold Copper and these other pesticides must not exceed 0.4 lb. a.i./acre/year. (5) Do not exceed the equivalent of 1.0 lb. a.i./A per crop of soil-applied mefenoxam-containing products at planting and 0.25 lb. a.i./A of foliar-applied mefenoxam-containing products. (6) Do not apply Ridomil Gold/Copper within 3 days prior to harvest (3-day PHI).

Ridomil Gold®, Bravo®, and the Syngenta logo are trademarks of a Syngenta Group Company

©2011 Syngenta

RID GOL COP 804 0211 SUPPLEM – pl – 6/7/11