



Section 2(ee) Recommendation

Date: June 27, 2014
Product: Ridomil Gold® SL
EPA Reg. No. 100-1202
Use: Addition of Late Blight (*Phytophthora infestans*) as a Pest Controlled on Potato
State: CT, NH, MA, ME, NJ, PA, RI, VT

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crop	Pest	Rate	Instructions
Potato	Late Blight (<i>Phytophthora infestans</i>)	3.2 oz/A (0.1 lb ai)	A tank mix with a labeled rate of mancozeb or chlorothalonil products is required. Begin preventative applications when conditions are favorable for disease. Apply on a 14 day schedule and alternate with an effective non-Group 4 fungicide. Use in conjunction with other management practices such as crop rotation and resistant varieties.

- Do not exceed the equivalent of 0.34 lb ai/A per season of soil-applied and 0.4 lb ai/A per season of foliar applied mefenoxam-containing products.
- Do not apply within 14 days of harvest (14-day PHI).

The user must refer to the federally approved labels and recently approved Supplemental Label, SCP 1202A-S5 0614, for the above noted product and read and follow all directions for use, restrictions, and precautions.

The user should have this recommendation in its possession at the time of use.

This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

Ridomil Gold® SL and the Syngenta logo are trademarks of a Syngenta Group Company.

©2014 Syngenta

2(ee) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Expiration Date: June 27, 2019

Label Code: CT1202014AC0614