

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP PRO® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Roundup PRO Herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.



EPA Reg. No. 524-475

Roundup PRO and Roundup Ready are registered trademarks of Monsanto Technology LLC.

TANK MIXTURES FOR IMPROVED CONTROL OF BENTGRASS (*Agrostis spp.*)

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY® CROPS) DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

USE INSTRUCTIONS

See "GENERAL INFORMATION", "MIXING" and "APPLICATION EQUIPMENT AND TECHNIQUES" sections of the label booklet for Roundup PRO herbicide for essential product performance information, and additional directions and restrictions on mixing and application of this product.

Roundup PRO herbicide may be tank mixed with the products listed below for improved control of bentgrass (*Agrostis spp.*).

When tank mixing, read and carefully observe label directions, cautionary statements and all information on the labels of each product used. Refer to each individual product label for approved use sites.

ENVOY®
FUSILADE® II
FUSION®
VANTAGE®

Dry ammonium sulfate, at 1 to 2 percent by weight, may be added to the spray solution for the above applications. The equivalent rate of ammonium sulfate in a liquid formulation may also be used. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

Broadcast Treatment

- Apply 2 to 4 quarts per acre of this product plus 34 fluid ounces per acre of **Envoy** in 20 to 40 gallons per acre of spray solution.
- Apply 2 to 4 quarts per acre of this product plus 24 fluid ounces per acre of **Fusilade II** in 20 to 40 gallons per acre of spray solution.

- Apply 2 to 4 quarts per acre of this product plus 60 fluid ounces per acre of **Vantage** in 20 to 40 gallons per acre of spray solution.
- Apply 2 to 4 quarts per acre of this product plus 9 fluid ounces per acre of **Fusion** in 20 to 40 gallons per acre of spray solution.

Re-treatment may be needed in the event of incomplete control.

Spot Treatment

- Mix 2.66 fluid ounces of this product with 1.3 fluid ounces of **Envoy** in one gallon of water and spray-to-wet.
- Mix 2.66 fluid ounces of this product with 0.75 fluid ounces of **Fusilade II** in one gallon of water and spray-to-wet.
- Mix 2.66 fluid ounces of this product with 3 fluid ounces of **Vantage** in one gallon of water and spray-to-wet.
- Mix 2.66 fluid ounces of this product with 0.25 fluid ounces of **Fusion** in one gallon of water and spray-to-wet.

ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Envoy is a registered trademark of Valent U.S.A. Corporation.

Fusilade and Fusion are registered trademarks of Syngenta Group Company.

Vantage is a registered trademark of BASF Corporation.

Read the “LIMIT OF WARRANTY AND LIABILITY” in the label booklet for Roundup PRO herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

©2004 MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167
3/18/04

21136C1-43