

GROW MORE®

GROW MORE®

GROW MORE®

GUARANTEED ANALYSIS
0.3 - 0.7 - 5

Total Nitrogen (N)0.30%
 0.30% Water Soluble Organic Nitrogen
 Available Phosphate (P₂O₅)0.70%
 Soluble Potash (K₂O)5.00%
 Derived from Seaweed *Ascophyllum Nodosum*
 5.1 Lbs. dry matter content per gallon
 Density 10.17 Lbs./Gal. (1.22 Kgs./Liter)

FOR COMMERCIAL USE ONLY

A NATURAL PRODUCT THAT LEAVES NO RESIDUE
 ON CROPS. MAY BE APPLIED ANYTIME DURING
 THE GROWING SEASON AND POST HARVEST.

CROPS	FOLIAR OR SOIL RATES AND TIMINGS
HORTICULTURE NURSERY/GREENHOUSE FOILAR APPLICATION & SOIL DRENCH	Use one (1) quart product per 250 gallons water or one (1) Liter Seaweed Extract in 1000 Liters water for a; 10 ppm rate
INJECTOR SYSTEM: Intermittent Feeding (1:100 Injector Ratio)	Add 3 pts (48 fl. ozs or 3 pts product in 100 gallons water in injector tank for 1:100 injection applications. (Metric = 1.5 Liters Seaweed Extract into 380 Liters water). Apply every 14 days throughout the growing season at 10 ppm. with fertilizers
Constant Feeding (1:100 Injector Ratio)	Add 8 to 16 fl. ozs Product per 100 gallons water Injection tank for 2-4 ppm with other fertilizers or pesticide products. (Metric: 250-500 ml into 380 Liters water)

Manufactured By
GROW MORE®
 15600 New Century Drive,
 Gardena, CA 90248 U.S.A.
www.growmore.com



0 180986 06091 5
 2 X 2.5 GALLONS

DIRECTIONS

For best results apply one pint (475 mL) in 100 gallons (380 Liters) of water per acre. Four correctly timed applications per growing season are recommended.

Apply as full coverage spray for optimum results.

Shake Before Using

Information regarding the contents and levels of metals in this product is available on the internet at:
<http://www.aapfco.org/metals.htm>

WARNING: This product contains chemicals known to the State of California to cause cancer and birth defects or other reproductive harm.

SEAWEED EXTRACT

Liquefied Natural Kelp



50%

COLD WATER PROCESSED
HELPS PROMOTES PLANT
GROWTH & VIGOROUS
ROOT SYSTEM
FOLIAR OR IRRIGATION



DENSITY: 10.17 LBS. PER GAL. (1.21Kg/Ltr)

NET WT.: ■ 2.5 Gallon ■ 55 Gallon ■ 275 Gallons

CROPS	FOLIAR OR SOIL RATES AND TIMINGS
FRUIT CROPS Stone Fruit, Cherries, Apple, Pears	(1.0-4.0 Pints product/Acre = 1.2 - 4.7 Liters/Ha) Pre-Bloom, Petal Fall, every 2-3 week until harvest, once post harvest for return bloom
Strawberries, Cane Berries, Blueberries	Transplant, Pre-Bloom, Fruit Set, every 7-14 days
Grapes: Table Grapes Wine, Juice, Raisin	8'- 12' cane, Pre-Bloom, Post-Bloom, every 14 days Pre-Bloom, Bloom, Post Bloom, Veraison, Post-harvest Seaweed Extract can increase fruit set; avoid bloom sprays in hard to thin crops like table grapes/apples.
Citrus (Orange, Lemon, Mandarin, etc.)	Spring Growth, before 1st Bloom, Petal Fall Summer, Fall, (Fresh Market 8 weeks before harvest)
NUT CROPS Almonds, Walnuts, all other nuts	(1.0-4.0 Pints product/Acre = 1.2 - 4.7 Liters/Ha) Pre-Bloom (bud stage), Petal Fall (Post-Bloom), Every 14 days (nut fill), Post Harvest for vigor
Avocados	Spring leaf growth, Pre-Bloom Petal Fall 14 day intervals during fruit sizing
VEGETABLE CROPS Tomatoes, Peppers, Eggplant	(1.0-4.0 Pints product/Acre = 1.2 - 4.7 Liters/Ha) Transplanting drench, 4 to 6" Growth. Pre. Bloom, 7-14 intervals through harvest
Cole Crops: Broccoli, Cauliflower, Kale, Cabbage, Collards, other greens	4-6 Leaf stage 14 days later head initiation, 14 day intervals for on going greens
Carrots, Onions, Leeks, Chives, Beans,	75% seedling emergence 3rd true leaf and root enlargement 14 day intervals, bulb swelling 4.6 True Leaf Stage Bud Stage and First Bloom, Pod Set 14 day intervals for sizing
Peas, Lentils, other legumes, Lettuce, Spinach, Celery, Cilantro, Parsley	7-14 day seedling 7 day intervals until harvest
FIELD CROPS: Small-Grain, Rice, Wheat, Oats, Sorghum, Canola, Chia, Quinoa	(1.0-4.0 Pints product/Acre = 1.2 - 4.7 Liters/Ha) 4'-6" seedling growth. First node. Flag leaf stage with fungicide applications
Potatoes, Sweet Potatoes Yams	75% emergence, Tuber initiation, Tuber set, Bloom time
Maize, Sweet Corn	4'- 6" growth, 10-14 growth, Just before tasseling
Soybeans	Early spring growth (3rd leaf), Full Bloom, 2 or more sprays on 2-3 week Intervals
Alfalfa Hay, Forage Crops. Pasture	Early Spring regrowth or establishment 14 day intervals or after cuttings.