



SUPPLEMENTAL LABEL

SONOMA 40WSP

READ THE ENTIRE LABEL FOR SONOMA 40WSP BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using SONOMA 40WSP as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

EPA Reg. No. 42750-141

AD092807

USE ON HOPS

KEEP OUT OF REACH OF CHILDREN

WARNING

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

For Use and Distribution Only in the States of Alaska, Idaho, Montana, Nevada, Oregon, Utah, Washington, and Wyoming)			
DISEASE	MYCLOBUTANIL 40WP AG (OZ/ACRE)	USE RECOMMENDATIONS	RESTRICTIONS
Powdery Mildew (Uncinula spp.)	2 – 4 (0.8 to 1.6 oz active)	Apply every 7 - 10 days during emergence to training growth stage. Do not apply less than 2 oz of SONOMA 40WSP per acre or adequate efficacy may not be achieved.	Do not make more than 4 applications of SONOMA 40WSP per acre per season. Do not apply more than 2.5 lb of SONOMA 40WSP (1 lb active ingredient) per acre per season. Pre-Harvest Interval: Do not apply within 14 days of harvest. Do not graze livestock in treated areas or harvest crops grown in treated areas for silage or hay. Chemigation: Do not apply through any type of irrigation system.
	4 – 6 (1.6 to 2.4 oz active)	Apply every 5 – 10 days during training to wire growth stage (prior to beginning of bloom when vines are rapidly growing). Do not apply less than 4 oz of SONOMA 40WSP per acre or adequate efficacy may not be achieved.	
	6 – 10 (2.4 to 4.0 oz active)	Apply every 7 – 10 days during wire to 14 day pre-harvest stage. Do not apply less than 6 oz of SONOMA 40WSP per acre or adequate efficacy may not be achieved.	
Apply SONOMA 40WSP in sufficient water for thorough coverage using ground equipment or by air in a minimum spray volume of 10 gallons of water per acre. Thorough coverage is essential. Use the shorter spray interval on susceptible varieties or under disease pressure. Resistance Management: SONOMA 40WSP belongs to the sterol demethylation inhibitor (DMI) class of fungicides and is classified as a Group 3 Fungicide by EPA. Since certain fungi can develop resistance to this class of products, the use of SONOMA 40WSP should be part of a resistance management strategy that includes alternation and/or tank mixing with fungicides of different modes of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.			

Manufactured By:
Albaugh, Inc.
1525 NE 36th Street
Ankeny, IA 50021