



TECHNICAL INFORMATION BULLETIN
FOR THE USE OF Stargus® BIOFUNGICIDE for
Berkeleyomyces (=Thielaviopsis) basicola (black root rot)
on Petunia

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(Ee) RECOMMENDATIONS ARE RECOGNIZED.

Stargus®

EPA Reg. No. 84059-28

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crop	Target Pathogen	Application Method	Product Use Rate per Application	Application Instructions
Petunia	<i>Berkeleyomyces</i> (= <i>Thielaviopsis</i>) <i>basicola</i> (black root rot)	Drench	1% v/v	Apply as a preventative drench to the growing media; can be repeated at weekly intervals as needed.
Stargus ® has a pre-harvest interval (PHI) of 0 days . Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours .				

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.

Stargus ® is a registered trademark of Marrone Bio Innovations, Inc.

© Copyright Marrone Bio Innovations, Inc., 2021

Marrone Bio Innovations, Inc.

1540 Drew Ave.

Davis, California 95618

Telephone (530) 750-2800

www.marronebio.com