

FIFRA Section 24(c) Special Local Need Label  
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON For  
use on Highbush Blueberries



EPA REG. NO. 34704-803

EPA SLN NO. OR-050016

This label valid until December 31, 2023 or until otherwise amended, withdrawn, canceled, or suspended.

ACTIVE INGREDIENT:

Dimethylamine salt of 2,4-Dichlorophenoxyacetic acid 46.6%

OTHER INGREDIENTS: 53.4%

TOTAL 100.0%

---

**KEEP OUT OF REACH OF CHILDREN**  
**DANGER PELIGRO**

---

**ENVIRONMENTAL HAZARDS**

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

**Groundwater Contamination:**

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This special local need label (SLN) label and the EPA-registered label must be in possession of the user at the time of pesticide application.

Follow all applicable directions, restrictions, and precautions on this SLN label and on the main EPA-registered label.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

**High bush blueberries (Vegetative strip between rows) / Broadleaf weeds (such as Broadleaf dock, Canada thistle, Dandelion, Spotted catsear):**

Apply 3 pints of Saber Herbicide (1.4 lb ae) in 50 gallons of water per acre using ground equipment only. The first application should be made in the spring as a direct, shielded spray to the vegetative strip between the blueberry rows, avoiding contact with the blueberry plant

foliage. For optimum herbicide performance, do not mow between rows for at least 7 days before or after the application. If necessary, a second application of Saber Herbicide at 3 pints (1.4 lb ae) in 50 gallons of water per acre may be made after harvest to control regrowth of broadleaf weeds.

**Restrictions:**

- Do not enter or allow worker entry during the restricted entry interval (REI) of 48 hours.
- A maximum of 3 pints (1.4 lb ae) of Saber Herbicide can be applied per acre per application.
- Do not make more than two applications of Saber Herbicide per acre per year.
- The Preharvest interval (PHI) for Saber Herbicide is 30 days.
- For use under this SLN label do not apply Saber Herbicide through any type of irrigation system.
- Do not apply Saber Herbicide by aerial application.
- Do not apply in or near greenhouses.
- Avoid applications of Saber Herbicide when winds are blowing toward nearby susceptible plants or when temperature inversions are expected. Do not make direct application or allow spray drift to contact susceptible plants since very small quantities of this herbicide can cause severe injury in the growing or dormant period. Plants contacted may be killed or suffer significant injury resulting in grade or yield losses.

**Precautions**

- Do not apply Saber Herbicide directly to blueberry plants or injury may result. To avoid injury to blueberry plants, do not apply if the temperature is 65 degrees or higher.
- Soil residue of this product may temporarily inhibit seed germination and plant growth.

24[c] Registrant:  
Loveland Products Inc.  
P.O. Box 1286  
Greeley, CO 80632-1286

Rev. 06/05/2019