

SATORI Fungicide

EPA REG NO. 34704-1068



FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the U.S. Environmental Protection Agency

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MICHIGAN, MINNESOTA, NORTH DAKOTA, MONTANA, WYOMING, NEBRASKA, COLORADO, IDAHO

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use on Sugar Beets

Target Diseases- Foliar	Use Rate Fluid ounces Satori/ Acre (lb ai/Acre)	Application Instructions
Alternaria Leaf Spot (<i>Alternaria</i> spp., <i>A. alternata</i>) Ascochyta Leaf Spot (<i>Ascochyta cynarae</i>) Rust (<i>Uromyces betae</i> , <i>Puccinia helianthi</i>) White Rust (<i>Albugo tragopogonis</i>)	6.0-20.0 (0.10-0.33)	For powdery mildew, make preventative applications on a 5- to 7- day schedule. For all other diseases, begin Satori Fungicide applications prior to disease development and continue throughout the season every 7-14 days. Follow resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.
Cercospora Leaf Spot (<i>Cercospora betae</i> , <i>C. pastinaceae</i>) Powdery Mildew (<i>Erysiphe polygoni</i> , <i>Leveillula taurica</i>)	9.0-15.5 (0.15-0.25)	Do not apply more than one application of Satori or other Group 11 fungicide before alternation with a fungicide not in Group 11.

Target Diseases- Soilborne	Use Rate fl oz/ 1000 row ft (lbs ai/ 1000 row ft)	Application Instructions
Circular Spot, Southern Blight (<i>Sclerotium rolfsii</i>) Pythium Root Rot (<i>Pythium aphanidermatum</i>) Rhizoctonia Stem Canker, Crown Rot (<i>Rhizoctonia solani</i>)	0.40-0.80 fl oz/ 1000 row ft (0.0065- 0.013 lb ai/1000 row ft)	For sugar beets apply 3–7-inch banded applications in a minimum of 10 gallons per acre at the 2- to 8-leaf stage. Do not apply as a dribble application over the seed row. Tank mixtures of Satori Fungicide with crop oil concentrates (COC) or methylated spray oil (MSO) may result in crop injury. If cool soil conditions are expected after planting which could result in an extended period of plant emergence, do not apply Satori Fungicide in-furrow. DO NOT use a starter fertilizer if using Satori Fungicide at the time of planting. NOTE: For soilborne/seedling disease control, see directions and rates under SOILBORNE/SEEDLING DISEASE CONTROL on Satori Fungicide Section 3 label.

Specific Use Restrictions:

- 1) **Maximum Single Application Rate:** Do not exceed the maximum rate listed in the table.
- 2) **Minimum Application Interval:** 5 days
- 3) **Maximum Annual Rate:** Do not apply more than 120 fl oz of product/A/year.
Do not apply more than 2.0 lb ai/A/year of azoxystrobin-containing products.
- 4) Do not apply more than 6 applications per year at the high rate (20.0 fl oz/A) or 20 applications per year at the low rate (6.0 fl oz/A). When applying at 9.0 fl oz/A, do not apply more than 13 applications per year. When applying at 15.5 fl oz/A, do not apply more than 7 applications per year.
- 5) Apply as an in-furrow spray in a minimum of 10 gallons per acre.
- 6) **Pre-Harvest Interval (PHI):** Satori Fungicide may be applied the day of harvest (0-day PHI).

Loveland Products, Inc.
P.O. Box 1286
Greeley, CO 80632