



Bayer CropScience LP
 800 N. Lindbergh Blvd.
 St. Louis, MO 63167
 1-866-99BAYER (1-866-992-2937)

SCALA® brand SC Fungicide

For suppression of Powdery Mildew in
 Grapes and Strawberry
 EPA Reg. No. 264-788

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF CALIFORNIA AND FLORIDA

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

| CROP | DISEASE SUPPRESSION | USE RATE | | USE INFORMATION |
|-------|---------------------|---|-------------------|---|
| | | FL OZ PER ACRE | LBS A.I. PER ACRE | |
| GRAPE | Powdery Mildew | SCALA BRAND SC FUNGICIDE Alone | | When using tank mix rates, add fungicide(s) registered for use against the targeted disease(s). Typically, first applications are made at early bloom, and/or berry touch to bunch closure, veraison, and pre-harvest. Use sufficient water to ensure penetration of the canopy and coverage of the flowers or bunches. RESTRICTIONS: Do not apply more than 36 fl oz/A (1.4 lbs a.i./A) of SCALA BRAND SC FUNGICIDE per crop. Do not apply within 7 days of harvest. |
| | | 18 | 0.70 | |
| | | SCALA BRAND SC FUNGICIDE In tank mixes | | |
| | | 9 - 18 | 0.35 - 0.70 | |

| CROP | DISEASE SUPPRESSION | USE RATE | | USE INFORMATION |
|------------|---------------------|---|-------------------|---|
| | | FL OZ PER ACRE | LBS A.I. PER ACRE | |
| STRAWBERRY | Powdery Mildew | SCALA BRAND SC FUNGICIDE Alone | | When using tank mix rates, add fungicide(s) registered for use against the targeted disease(s). For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. Make applications from pre-bloom to harvest on a 7 to 14 day interval depending upon disease conditions. Always follow resistance management directions listed on this label and the recommendations for your area. RESTRICTIONS: Do not apply more than 54 fl oz/A (2.1 lbs a.i./A) of SCALA BRAND SC FUNGICIDE per crop. Do not apply within 1 day of harvest. |
| | | 18 | 0.70 | |
| | | SCALA BRAND SC FUNGICIDE In tank mixes | | |
| | | 9 - 18 | 0.35 - 0.70 | |

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*