



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

SCALA® BRAND SC Fungicide

EPA Reg. No. 264-788

For Use On: Almonds, Pistachios, Bulb Vegetables, Grapes, Stone Fruits (Except Cherries), Pome Fruits, and Strawberries

This supplemental label expires January 19, 2015 and must not be used or distributed after this date.

Supplemental Label

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Read this label and the product package label before using this product. This Supplemental Label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the product label for SCALA BRAND SC Fungicide attached to the container.

USE DIRECTIONS FOR SPECIFIC CROPS

ALMONDS, PISTACHIOS

CROP	DISEASE	USE RATE		USE INFORMATION
		FL. OZ. PER ACRE	LBS. A.I. PER ACRE	
Almond Pistachio	Brown rot blossom blight <i>(Monilinia laxa, M. fructicola)</i> Jacket rot, Green fruit rot (on almonds) <i>(Botrytis cinerea)</i>	SCALA BRAND SC FUNGICIDE Alone		For optimum results, begin applications as soon as crop and/or environmental conditions become favorable for disease development. Make applications on a minimum 7 day interval depending upon disease conditions. Do not apply more than 54 fl. oz./A (2.1 lbs. a.i./A) of SCALA BRAND SC FUNGICIDE per crop. Do not apply within 30 days of harvest. When using tank mix rates, add fungicide(s) registered for use against the targeted disease(s).
		9 -18	0.35 - 0.70	
	SCALA BRAND SC FUNGICIDE Alone			
	18	0.70		
	Shot hole <i>(Wilsonomyces carpophilus)</i> Botryosphaeria blight <i>(Botryosphaeria dothidia)</i> Scab (Suppression only) <i>(Cladosporium spp.)</i> Blossom & shoot blight (on pistachios) <i>(Botrytis cinerea)</i>	SCALA BRAND SC FUNGICIDE In tank mixes		
		9-18	0.35-0.70	

BULB VEGETABLES

CROP	DISEASE	USE RATE		USE INFORMATION
		FL. OZ. PER ACRE	LBS. A.I. PER ACRE	
Onion Green Dry Bulb Welsh Garlic including Great-Headed Leek Shallot	Botrytis leaf blight, Neck rot (<i>B. squamosa</i> <i>B. allii</i>)	SCALA BRAND SC FUNGICIDE Alone		When using tank mix rates, add fungicide(s) registered for use against the targeted disease(s). Use lower rate only in a tank mix with a broad spectrum fungicide for bulb vegetables. For optimum results, begin applications as soon as crop and/or environmental conditions become favorable for disease development. Make applications on a 7 to 14 day interval depending upon disease conditions. Do not apply more than 54 fl. oz./A (2.1 lbs. a.i./A) of SCALA BRAND SC FUNGICIDE per crop. Do not apply within 7 days of harvest.
		18	0.70	
	Purple blotch (<i>Alternaria porri</i>)	SCALA BRAND SC FUNGICIDE In tank mixes		
9-18		0.35-0.70		

GRAPES

CROP	DISEASE	USE RATE		USE INFORMATION
		FL. OZ. PER ACRE	LBS. A.I. PER ACRE	
Grape	Gray mold (<i>Botrytis cinerea</i>)	SCALA BRAND SC FUNGICIDE Alone		When using tank mix rates, add fungicide(s) registered for use against the targeted disease(s). Use lower rate only in a tank mix with another fungicide active against <i>Botrytis</i> . Apply product at the critical timings for <i>Botrytis</i> control. Typically, first applications are made at early bloom, and/or berry touch to bunch closure, veraison, and pre-harvest. Use sufficient water to ensure penetration of the canopy and coverage of the flowers or bunches. Do not apply more than 36 fl. oz./A (1.4 lbs. a.i./A) of SCALA BRAND SC FUNGICIDE per crop. Do not apply within 7 days of harvest.
		18	0.70	
		SCALA BRAND SC FUNGICIDE In tank mixes		
		9-18	0.35-0.70	

**STONE FRUITS
(Except Cherries)**

CROP	DISEASE	USE RATE		USE INFORMATION
		FL. OZ. PER ACRE	LBS. A.I. PER ACRE	
Apricot Nectarine Peach Plum (All varieties) Plum, Chickasaw Plum, Damson Plum, Japanese Plum, Stanley Plumcot Prune	Brown rot blossom blight (<i>Monilinia</i> spp.)	SCALA BRAND SC FUNGICIDE Alone		For optimum results, begin applications as soon as crop and/or environmental conditions become favorable for disease development. Apply when bud tissue is susceptible to infection (i.e., pink, white or red bud). If conditions favorable for disease development persist or recur, apply at full bloom or at petal fall. Make pre-harvest applications on 7 day intervals or longer. Do not apply more than three applications of SCALA BRAND SC FUNGICIDE alone or more than 54 fl. oz./A (2.1 lbs. a.i./A) of SCALA BRAND SC FUNGICIDE per crop. Do not apply within 2 days of fruit harvest or make more than 2 applications of a Group 9 fungicide within 30 days of harvest. Do not use on cherries. When using tank mix rates, add fungicide(s) registered for use against the targeted disease(s).
		9 - 18	35 – 0.70	
	Jacket rot, Green fruit rot (<i>Botrytis cinerea</i>)	SCALA BRAND SC FUNGICIDE Alone		
		18	0.70	
	Shot hole (<i>Wilsonomyces carpophilus</i>)	SCALA BRAND SC FUNGICIDE In Tank Mixes		
		9-18	0.35-0.70	
Brown fruit rot (<i>Monilinia fructicola</i>)				

POME FRUITS

CROP	DISEASE	USE RATE		USE INFORMATION
		FL. OZ. PER ACRE	LBS. A.I. PER ACRE	
Apple Pear (All varieties) Pear, Oriental Crabapple Loquat Mayhaw Quince	Scab (<i>Venturia</i> spp.)	SCALA BRAND SC FUNGICIDE Alone		Under conditions favorable for disease development, shorten the spray intervals and/or use the high rate. When using tank mix rates, add fungicide(s) registered for use against the targeted disease(s). For optimum results, begin applications at green tip or as soon as crop and/or environmental conditions become favorable for disease development. Make pre-harvest applications on 7 day intervals or longer. Do not apply more than 40 fl. oz./A (1.6 lbs. a.i./A) of SCALA BRAND SC FUNGICIDE per crop. Do not apply within 72 days of harvest.
		7 - 10	0.27 - 0.39	
		SCALA BRAND SC FUNGICIDE In tank mixes		
		5-10	0.19-0.39	

STRAWBERRIES

CROP	DISEASE	USE RATE		USE INFORMATION
		FL. OZ. PER ACRE	LBS. A.I. PER ACRE	
Strawberry	Gray mold (<i>Botrytis cinerea</i>)	SCALA BRAND SC FUNGICIDE Alone		<p>When using tank mix rates, add fungicide(s) registered for use against the targeted disease(s).</p> <p>For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. Make applications from pre-bloom to harvest on a 7 to 14 day interval depending upon disease conditions.</p> <p>Do not apply more than 54 fl. oz./A (2.1 lbs. a.i./A) of SCALA BRAND SC FUNGICIDE per crop. Always follow resistance management directions listed on this label and the recommendations for your area. Do not apply within 1 day of harvest.</p>
		18	0.70	
		SCALA BRAND SC FUNGICIDE In tank mixes		
		9-18	0.35-0.70	

**For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)**

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).