



Supplemental Labeling

AMVAC Scepter 70DG Herbicide

EPA Reg. No. 241-306-5481

**FOR USE IN EASTERN COTTONWOOD (*Populus deltoids*) AND HYBRID POPLAR CLONES INCLUDING
CROSSES BETWEEN *Populus deltoids*, *Populus Maximowiczii*, *Populus nigra* and *Populus trichocarpa*
GROWN ON FIBER FARMS**

ONLY FOR DISTRIBUTION AND USE IN THE FOLLOWING STATES

Alabama	Kansas	Missouri	South Carolina
Arkansas	Kentucky	Nebraska	South Dakota
Florida	Louisiana	New Jersey	Tennessee
Georgia	Maine	North Carolina	Texas
Idaho	Maryland	Ohio	Virginia
Illinois	Michigan	Oklahoma	West Virginia
Indiana	Minnesota	Oregon	Washington
Iowa	Mississippi	Pennsylvania	Wisconsin

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE SCEPTER[®] 70 DG HERBICIDE LABELING BEFORE USING. SEE THE SCEPTER 70 DG LEAFLET LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION, WEEDS CONTROLLED, USES WITH OTHER PRODUCTS AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. The use of SCEPTER 70 DG not consistent with this labeling may result in injury to crops. DO NOT apply this product through any type of irrigation system.

This labeling must be in the possession of the user at the time of herbicide application.

GENERAL INFORMATION

SCEPTER 70 DG herbicide applied preplant incorporated, preemergence or postemergence provides early season weed control and residual activity on broadleaf weeds and grass weeds in eastern cottonwood and hybrid poplars, grown on fiber farms or tree plantations.

When tank-mixing herbicides, always follow the most restrictive label.

APPLICATION INSTRUCTIONS

Applications prior to planting trees: Apply 2.8 to 5.6 oz SCEPTER 70 DG per acre preplant incorporated, preemergence or postemergence to control labeled weeds. For any of these methods tank mix with 1.2 to 4.8 quarts PENDULUM[®] 3.3 EC herbicide to improve preemergence control of weeds listed on the PENDULUM 3.3 EC label.

For preplant incorporated applications, SCEPTER 70 DG may be tank mixed with trifluralin. Refer to the trifluralin label for specific use recommendations, rates and weeds controlled.

For postemergence applications, glyphosate may be used with SCEPTER 70 DG alone or in combination with PENDULUM 3.3 EC. Refer to the glyphosate label for specific use recommendations, rates and weeds controlled.

Postplant applications prior to trees breaking dormancy (bud swell stage): Apply 2.8 to 5.6 oz SCEPTER 70 DG per acre preemergence or postemergence to control labeled weeds. Tank mix with 1.2 to 4.8 quarts PENDULUM 3.3 EC to improve preemergence control of weeds listed on the PENDULUM 3.3 EC label.

Postplant applications on actively growing trees: Apply 2.8 to 5.6 oz SCEPTER 70 DG per acre preemergence or postemergence to control labeled weeds. Different clonal lines may vary in their tolerance to SCEPTER 70 DG. Therefore, before making over the top applications on actively growing trees, spray a few trees and observe for plant damage over a period of six weeks prior to full-scale application. For best crop safety on actively growing trees, do not include a surfactant. Do not tank mix with PENDULUM 3.3 EC herbicide.

Do not apply more than 5.6 oz SCEPTER 70 DG per acre per year.

Refer to the SCEPTER 70 DG label for additional information on application instructions.

WEEDS CONTROLLED

When applied as directed, SCEPTER 70 DG herbicide will provide postemergence control of the following weed species. Refer to SCEPTER 70 DG label for weeds controlled by soil applications.

5.6 oz. SCEPTER 70 DG per acre

<u>Common Name</u>	<u>Scientific Name</u>
Broadleaf Weeds	
Carpetweed	Mollugo verticillata
Cocklebur, common	Xanthium strumarium
Eclipta	Eclipta prostrata
Jimsonweed	Datura stramonium
Mustard, wild	Brassica kaber
Nightshade, eastern black	Solanum ptycanthum
Poinsettia, wild	Euphorbia heterophylla
Prickly sida	Sida spinose
Smartweed, ladysthumb	Polygonum persicaria
Smartweed, Pennsylvania	Polygonum pennsylvanicum
Sunflower, common	Helianthus annuus
Pigweed	
Palmer	Amarathus palmeri
Redroot	Amaranthus retroflexus
Smooth	Amaranthus hybridus
Spiny	Amaranthus spinosus
Venice mallow	Hibiscus trionum
Grasses	
Corn, volunteer*	Zea mays
Foxtail, giant	Setaria faberi
Foxtail, yellow	Setaria glauca
Panicum, fall	Panicum dichotomiflorum

* A postemergence application of SCEPTER 70 DG will control only those field corn hybrids which DO NOT possess tolerance or resistance to SCEPTER 70 DG or other imidazolinone herbicides.

Scepter and Pendulum are registered trademarks of BASF.



AMVAC Chemical Corporation
4100 E. Washington Blvd.
Los Angeles, CA 90023 U.S.A.
1-888-462-6822
20151222