

Product Bulletin

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Sequoia™

EPA Reg. No. 62719-623

Special 2(ee) Recommendation†

(For Distribution and Use in the States of Arizona and Hawaii)

For the Control of Western Tarnished Plant Bug (Lygus Bug) on Specific Crops

Refer to product label for Sequoia for general use precautions, mixing directions, application directions, and restrictions.

ATTENTION

†This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Sequoia insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Sequoia according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Sequoia insecticide.

Directions for Use

Apply Sequoia to Brassica Head and Stem Vegetables (Crop Group 5-16) and Kolrabi, Leafy Greens (Subgroup 4-16A) Leafy Petiole Vegetable (Subgroup 22B), Celtuce, Florence Fennel, and Watercress at the rate of 4.25 to 5.75 fl. oz/acre to control Western Tarnished Plant Bug (Lygus bug). Use a higher rate in the rate range for heavy pest populations.

Restrictions:

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 7 days apart.
- Do not make more than four applications per crop.
- Do not make more than two consecutive applications per crop.
- Do not apply more than a total of 17 fl oz of Sequoia (0.266 lb ai of sulfoxaflor) per acre per year.
- Do not apply this product at any time between 3 days prior to bloom and until after petal fall.

™® Trademarks of Corteva Agriscience and its affiliated companies

R919-023

Issued: 08/17/22

Replaces: R919-022