

Product Bulletin

Corteva Agriscience

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Sequoia[®] CA

EPA Reg. No. 62719-728

Special 2(ee) Recommendation[†] (For Distribution and Use in the State of California)

For the Control of Glassy-winged Sharpshooter in Grapes

Refer to product label for Sequoia CA for general use precautions, mixing directions, application directions, and restrictions.

ATTENTION

[†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Sequoia CA insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Sequoia CA according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Sequoia CA.

Directions for Use

Apply Sequoia CA to Grapes at the rate of 4.5 to 5.75 fl. oz/acre to control Glassy-winged Sharpshooter. Use a higher rate in the rate range for heavy pest populations.

Restrictions:

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 7 days apart.
- Do not make more than four applications per crop.
- Do not make more than two consecutive applications per crop.
- Do not apply more than a total of 17 fl oz of Sequoia CA (0.266 lb ai of sulfoxaflor) per acre per year.
- Do not apply this product until after petal fall.
- If blooming vegetation is present 12 feet out from the downwind edge of the field, a downwind 12-foot on-field buffer must be observed.

[™][®] Trademarks of Corteva Agriscience and its affiliated companies

R447-036

Issued: 03/10/2021

Initial printing.