



Recommendation

FIFRA SECTION 2(ee) RECOMMENDATION

GROUP	28	INSECTICIDE
-------	----	-------------

Shenzi 400 SC Insecticide

(EPA Reg. No. 70506-607)

REDUCED RATES FOR CONTROL OF BEET ARMYWORM AND WESTERN YELLOWSTRIPPED ARMYWORM IN ALFALFA IN STATES OF ARIZONA, CALIFORNIA, OREGON AND WASHINGTON

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by USEPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all applicable instructions, restrictions, and precautions, and Conditions of Sale and Warranty on the EPA-approved label. This recommendation and the federally registered label must be in the possession of the user at the time of pesticide application.

CROP	APPLICATION METHOD	INSECT	SHENZI 400 SC INSECTICIDE RATE PER ACRE IN FL OZ (LB A.I.)	RE-TREATMENT INTERVAL IN DAYS (RTI)	PRE-HARVEST INTERVAL IN DAYS (PHI)	RE-ENTRY INTERVAL IN HOURS (REI)
Alfalfa	Foliar	Beet Armyworm Yellowstriped Armyworm	1.3 – 2.5 (0.034 – 0.065)	n/a	0	4

APPLICATION INSTRUCTIONS:

SHENZI SC INSECTICIDE can be applied by overhead sprinkler chemigation systems. See CHEMIGATION USING OVERHEAD SPRINKLER SYSTEMS section for instructions on overhead sprinkler chemigation.

RESTRICTIONS:

DO NOT make more than 4 applications per acre per calendar year.

DO NOT apply more than 7.7 fl oz of SHENZI SC INSECTICIDE or 0.2 lb a.i. chlorantraniliprole containing products per acre per calendar year. Apply higher rates within the listed range for heavier infestations, larger/denser crops, or extreme environmental conditions such as rainy weather and high temperatures. Make one application per cutting.

Do not make more than two sequential applications of SHENZI SC INSECTICIDE before rotating to another registered insecticide having a different mode-of-action.