

Technical Bulletin Signaling

Foliar Application

Crops	Rate	Application Timing
Vegetable Crops: Leafy Greens, Cole Crops, Row Crops, Pepper, Tomato, etc.	24-32 oz per acre	At planting and every 2 or 3 weeks on development, flowering and bearing stages
Fruit Crops: Avocado, Citrus, Nectarine, Blackberry, Blueberry, Raspberry, Strawberry Table Grapes, Wine Grapes, etc.	28-36 oz per acre	On nursery every 3 or 4 weeks On nursery every 3 or 4 weeks. At planting or season beginning and every 2 or 3 weeks on development, flowering and bearing stages On nursery every 3 or 4 weeks- At planting or season beginning twice or three times 3 weeks after
Nut Crops: Almond, Pistachio, Walnut, etc.	28-36 oz per acre	On nursery every 3 or 4 weeks
Field Crops: Carrot, Corn, Cotton, onion, Potato, Rice, etc.	24-32 oz per acre	As preplant treatment or seeding or and every 2 or 3 weeks on development, flowering and bearing stages
Forage Crops: Alfalfa, Fescue, Bluegrass, etc.	24-32 oz per acre	At seeding and once or twice 15 days after. Once after each harvest.

Drip Applications

Crops	Rate	Application Timing
Vegetable Crops: Leafy Greens, Cole Crops, Row Crops, Pepper, Tomato, etc.	24-36 oz per acre	At planting and every 2 or 3 weeks on development, flowering and bearing stages
Fruit Crops: Avocado, Citrus, Nectarine, Pomegranate, Blackberry, Blueberry, Raspberry, Strawberry	28-36 oz per acre	On nursery every 3 or 4 weeks On nursery every 3 or 4 weeks. At planting or season beginning and every 2 or 3 weeks on development, flowering and bearing stages

Table Grapes, Wine Grapes, etc.		On nursery every 3 or 4 weeks- At planting or season beginning twice or three times 3 weeks after
Nut Crops: Almond, Pistachio, Walnut, etc.	28-36 oz per acre	On nursery every 3 or 4 weeks
Field Crops: Carrot, Corn, onion, Potato, Cotton, Rice, etc.	20-32 oz per acre	As preplant treatment or seeding or and every 2 or 3 weeks on development, flowering and bearing stages
Forage Crops: Alfalfa, Fescue, Bluegrass, etc.	20-32 oz per acre	At seeding and once or twice 15 days after. Once after each harvest.

Signaling is not limited for the crops listed above it may be use with any type of plant.



1444 Fulton Fresno CA 93721